

Time variability and heterogeneity in the coma of 67P/Churyumov-Gerasimenko

M. Hässig,^{1,2*} K. Altwegg,^{1,3} H. Balsiger,¹ A. Bar-Nun,⁴ J. J. Berthelier,⁵ A. Bieler,^{1,6} P. Bochsler,¹ C. Briois,⁷ U. Calmonte,¹ M. Combi,⁶ J. De Keyser,^{8,9} P. Eberhardt,^{1†} B. Fiethe,¹⁰ S. A. Fuselier,² M. Galand,¹¹ S. Gasc,¹ T. I. Gombosi,⁶ K. C. Hansen,⁶ A. Jäckel,¹ H. U. Keller,^{12,14} E. Kopp,¹ A. Korth,¹³ E. Kührt,¹⁴ L. Le Roy,³ U. Mall,¹³ B. Marty,¹⁵ O. Mousis,¹⁶ E. Neefs,¹⁷ T. Owen,¹⁸ H. Rème,^{19,20} M. Rubin,¹ T. Sémon,¹ C. Tornow,¹⁴ C.-Y. Tzou,¹ J. H. Waite,² P. Wurz¹

Comets contain the best-preserved material from the beginning of our planetary system. Their nuclei and comae composition reveal clues about physical and chemical conditions during the early solar system when comets formed. ROSINA (Rosetta Orbiter Spectrometer for Ion and Neutral Analysis) onboard the Rosetta spacecraft has measured the coma composition of comet 67P/Churyumov-Gerasimenko with well-sampled time resolution per rotation. Measurements were made over many comet rotation periods and a wide range of latitudes. These measurements show large fluctuations in composition in a heterogeneous coma that has diurnal and possibly seasonal variations in the major outgassing species: water, carbon monoxide, and carbon dioxide. These results indicate a complex coma-nucleus relationship where seasonal variations may be driven by temperature differences just below the comet surface.

Initially, comets were classified depending on the location where they formed in the protoplanetary disc (1, 2). This classification assumed a similar composition of the nucleus within a given formation region. No come-

tary nucleus composition has been sampled in situ. Rather, it is implicitly assumed that measurements of the outgassing of comets reveal the composition of the volatile components of the nucleus. However, compositional homogeneity of at least one comet was confirmed by studying outgassing from the fragments of the broken-up comet Schwassmann-Wachmann 3 (3). Detailed observations of other cometary comae indicated that there is evidence of heterogeneity. Missions

to comet Halley detected release of volatiles in multiple jet-like features that were dominantly seen on the sunlit side of the nucleus (4, 5). The Deep Impact mission detected asymmetries in composition in the coma of Tempel 1 (6). In particular, these remote sensing observations at Tempel 1 indicated an absence of correlation between H₂O and CO₂ in the coma.

Detailed, close-up cometary images have also showed visible differences between different areas of cometary nuclei. These images suggested that heterogeneity in the coma of a comet may be related to heterogeneity of the nucleus. Observations by EPOXI (Extrasolar Planet Observation and Deep Impact Extended Investigation) at Hartley 2 in 2010 near perihelion indicated that the nucleus is complex, with two different sized lobes separated by a middle waist region that is smoother and lighter in color (7). Outgassing from sunlit surfaces of the nucleus revealed that the waist and one of the lobes were very active. A CO₂ source was detected at the small lobe of the comet, whereas the waist was more active in H₂O and had a considerably lower CO₂ content. Based on these coma observations, it has been tentatively suggested that the heterogeneity in the comet's nucleus was primordial (7). Seasonal effects could not be ruled out because the observations also showed a complex rotational state for the comet (7). The smaller of the two lobes may have been illuminated differently because of this complex rotation (7). In support of the findings at Hartley 2, there are indications of a heterogeneous nucleus for comet Tuttle and a heterogeneous coma (7, 8).

The Stardust mission to comet P81/Wild 2, on the other hand, showed a large mixing of materials on the scale of grains and therefore a homogenized mix of the refractory material in the comet (9). The results at Hartley 2 and at

¹Physikalisches Institut, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland. ²Southwest Research Institute, 6220 Culebra Road, San Antonio, TX 78238, USA. ³Center for Space and Habitability (CSH), University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland. ⁴Department of Geosciences, Tel-Aviv University, Ramat-Aviv, Tel-Aviv, Israel. ⁵Laboratoire Atmosphères, Milieux, Observations Spatiales (LATMOS), Institut Pierre Simon Laplace (IPSL), Centre national de recherche scientifique (CNRS), Université Pierre et Marie Curie (UPMC), Université de Versailles Saint-Quentin-en-Yvelines (UVSQ), BP 102, UPMC, 4 Place Jussieu, F-75252 Paris Cedex 05, France. ⁶Department of Atmospheric, Oceanic and Space Sciences, University of Michigan, 2455 Hayward Street, Ann Arbor, MI 48109, USA. ⁷Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), UMR 7328 CNRS – Université d'Orléans, France. ⁸Belgian Institute for Space Aeronomy (BIRA-IASB), Ringlaan 3, B-1180 Brussels, Belgium. ⁹Center for Plasma Astrophysics, KU Leuven, Celestijnenlaan 200D, 3001 Heverlee, Belgium. ¹⁰Institute of Computer and Network Engineering (IDA), TU Braunschweig, Hans-Sommer-Strasse 66, D-38106 Braunschweig, Germany. ¹¹Department of Physics, Imperial College London, Prince Consort Road, London SW7 2AZ, UK. ¹²Institute for Geophysics and Extraterrestrial Physics, Technische Universität (TU) Braunschweig, 38106 Braunschweig, Germany. ¹³Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany. ¹⁴German Aerospace Center, Institute of Planetary Research, Asteroids and Comets, Rutherfordstrasse 2, 12489 Berlin, Germany. ¹⁵Centre de Recherches Pétrographiques et Géochimiques (CRPG), 15 Rue Notre Dame des Pauvres, BP 20, 54501 Vandœuvre lès Nancy, France. ¹⁶Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille), UMR 7326, 13388, Marseille, France. ¹⁷Engineering Division, BIRA-IASB, Ringlaan 3, B-1180 Brussels, Belgium. ¹⁸Institute for Astronomy, University of Hawaii, Honolulu, HI 96822, USA. ¹⁹Université de Toulouse, Université Paul Sabatier (UPS), Observatoire de Midi-Pyrénées (OMP), Institut de Recherche en Astrophysique et Planétologie (IRAP), Toulouse, France. ²⁰CNRS, IRAP, 9 Avenue du Colonel Roche, BP 44346, F-31028 Toulouse Cedex 4, France. *Corresponding author. E-Mail: myrtha.haessig@swri.org †Deceased.




Fig. 1. H₂O, CO, and CO₂ measurements for 4 to 8 August 2014. The upper panel shows the signal on the DFMS detector for H₂O, CO, and CO₂, and the lower panel shows the latitude and longitude of the nadir view of the spacecraft. At the top is the distance from the spacecraft to the comet. The signal increases with decreasing distance to the comet, and diurnal variations are also visible. CO₂ has a different periodicity than H₂O, as seen around 4 to 6 August.

P81/Wild 2 raise the larger question of whether heterogeneity in the coma is a common feature in comets and whether this reveals an underlying heterogeneity in the composition of the nucleus, which would point to general transport of cometesimals in the early solar system.

In August, the European Space Agency's mission Rosetta arrived at its target comet 67P/Churyumov-Gerasimenko (67P) after a 10-year journey (10). Rosetta provides an excellent opportunity for long-term study during the comet's sunward approach to perihelion. The observations presented here are from a 2-month period beginning near the initial encounter at about 3.5 astronomical units from the Sun.

Like Hartley 2, the nucleus of 67P appears complex in shape. The comet 67P consists of two lobes of different sizes, connected by a neck region. The lobes are much larger, more rugged, and darker than the neck region and the overall shape has been compared to a rubber duck (11). The structural similarities of 67P and Hartley 2 suggest the possibility of another heterogeneous comet and, by virtue of the extended observations at 67P, a chance to determine whether heterogeneity in the coma and nucleus are related.

Here, we show compositional variations in H₂O, CO, and CO₂ at comet 67P observed with ROSINA/DFMS (Rosetta Orbiter Spectrometer for Ion and Neutral Analysis/Double Focusing Mass Spectrometer) (12). ROSINA/DFMS is a mass spectrometer that measures the in situ neutral and plasma coma composition at the position of the spacecraft (see the supplementary materials). During Rosetta's approach to 67P, ROSINA/DFMS measured the neutral coma composition with a time resolution (>10 measurements per rotation) much finer than the rotation period of the comet of ~12.4 hours (13). In August, the spacecraft scanned the comet at northern summer hemisphere (positive latitudes) from about 10° up to almost 90° (Coordinates: Cheops System). In September, the spacecraft made a similar scan at southern winter hemisphere (negative latitudes) down to about -50°. Two data sets are shown in Figs. 1 and 2 to illustrate the diurnal and latitudinal variations and heterogeneity of the cometary coma.

During this approach and latitude scan, the H₂O, CO, and CO₂ signals from the comet increased by more than an order of magnitude, roughly in agreement with a $1/R^2$ dependence on the coma density, where R is the distance from the comet's center. Overall, the H₂O signal is the strongest; however, there are clearly periods when the CO or CO₂ signals rival that of H₂O.

Superposed on this general increase in signal are large, diurnal variations for all three neutral species. For H₂O, these variations are periodic, initially with half the rotation rate of the comet (~6.2 hours) and then, after 6 August, at the rotation rate (~12.4 hours). This change in periodicity in the signal is interpreted as a latitudinal effect of the sampling position. Peaks occur at $\pm 90^\circ$ longitude. For the most part, the CO signal follows the H₂O signal, but the variations are smaller. CO₂ shows a different periodicity. Initially, a CO₂ peak is observed in association with

an H₂O peak, and a second CO₂ peak occurs approximately 3 hours later. After August 6, a single CO₂ peak is observed; however, this peak is not exactly coincident with the H₂O peak. The two CO₂ peaks merge, resulting in a shoulder on the main peak and a slight shift of the main CO₂

peak relative to that of H₂O (~45 min or one measurement point). Statistical uncertainties ($\sqrt{\text{#particles}}$) in the signal detected by ROSINA/DFMS are smaller than the dots in Figs. 1 to 3, and contributions to the signal due to spacecraft outgassing (14) are subtracted.




Fig. 2. H₂O, CO, and CO₂ measurements for 15 to 19 September 2014. Over this 4-day period, the spacecraft remained at a nearly fixed distance from the comet and executed a southern latitude scan from about 0° to -45° latitude. H₂O and CO₂ have different periodicities, and there are deep minima in the H₂O signal. CO follows the CO₂ profile with less variation.




Fig. 3. H₂O, CO, and CO₂ profiles for 18 September 2014. The Sun is shining on the comet from the top middle of the pictures. The snapshots of the spacecraft view of the comet show that H₂O peaks are observed when the neck region is in view. The separate CO₂ peak and the deep minimum in H₂O occur when the spacecraft views the larger of the two lobes and the neck region is blocked. [Shape model credit: ESA/Rosetta/MPS for OSIRIS Team MPS/UPD/LAM/IAA/SSO/INTA/UPM/DASP/IDA].

The diurnal variations at half the rotation rate of the comet that are seen in August are also observed at southern latitudes in the September time frame (Fig. 2). The H₂O peaks in Fig. 3 are nearly equal, and there is a deep minimum between the two peaks. As in the first data set, CO follows H₂O. However, there is much less variation in CO than in H₂O, resulting in times when the CO signal is greater than that for H₂O. The best example of the differences between H₂O and CO₂ are seen just after 18 September (Figs. 2 and 3). The nearly equal H₂O peaks and the deep minimum in the H₂O signal are evident, as is the clear offset between the second CO₂ and H₂O peaks.

We have combined the signal and the spacecraft perspective over the 18 to 19 September 2014 window to illustrate which side of the comet is in view when the peaks occur (Fig. 3). The peaks in H₂O signal are observed when the neck of the comet is in view of the spacecraft. The deep minimum in H₂O signal is observed when the spacecraft views the southern hemisphere of the larger of the two lobes. This large lobe blocks a direct view of the neck of the comet. The separate, second CO₂ enhancement is observed when the spacecraft views the underside of the body of the larger of the two lobes of the comet. The CO signal in the second rotation of the comet follows the CO₂ profile, and CO and CO₂ have very similar intensities.

We see from this data (Figs. 1 to 3) that the coma composition of 67P is highly heterogeneous. H₂O, CO, and CO₂ variations are strongly tied to the rotation period of the comet and to the observing latitude. At large negative latitudes, the H₂O signal varies by at least two orders of magnitude (Fig. 3). Also, the H₂O minima are not as deep when the spacecraft is at mid and high positive latitudes because there is a view of the neck region over the edge of the larger lobe (see Fig. 1 and the observations on 15 September in Fig. 2).

The separate CO₂ peak also occurs when the spacecraft views the bottom of the larger of the two lobes of the comet (see Fig. 3 at 5 hours). CO follows H₂O at positive latitudes and follows both H₂O and CO₂ at negative latitudes.

The separate CO₂ peak, the large variations in the H₂O signal, and the weaker variations in CO result in large changes in the relative concentration of H₂O, CO, and CO₂ in the heterogeneous coma of 67P (see the supplementary materials). For example, the CO/H₂O number density ratio is 0.13 ± 0.07 , and the CO₂/H₂O ratio is 0.08 ± 0.05 in the last H₂O peak on 7 August at 18 hours in Fig. 1 (measured high in the northern summer hemisphere). However, the CO/H₂O ratio changes from 0.56 ± 0.15 to 4 ± 1 and back to 0.38 ± 0.15 within the second cometary rotation (Fig. 3), between 12 and 24 hours on 18 September, measured low in the southern winter hemisphere. Similarly, the CO₂/H₂O ratio changes from 0.67 ± 0.15 to 8 ± 2 and back to 0.39 ± 0.15 over the same rotation. These are large changes within a short amount of time, which indicate a strongly heterogeneous and time-variable coma.

The similarities in the structure of the nuclei and the heterogeneous comae of 67P and Hartley 2 are striking. The behavior in terms of the H₂O dominant outgassing at the neck versus CO₂ outgassing at one of the lobes described here was also found for Hartley 2 (7).

The compositional differences in the Hartley 2 coma were interpreted as evidence for a heterogeneous cometary nucleus (7). However, seasonal effects could not be ruled out. With observations over a wide range of latitudes at 67P, we can distinguish between compositional differences and seasonal effects; to do so, we have mapped the CO₂/H₂O density ratio from 17 August through 22 September onto the shape model (Fig. 4).

Although a direct mapping of the signal observed in the coma onto the comet surface is

oversimplified, a generalized interpretation reveals features of the outgassing of the comet. Seasonal effects on the CO₂/H₂O ratio are clearly evident (Fig. 4). On the upper half of the comet, the CO₂/H₂O ratio is less than 1, indicating a higher sublimation of H₂O from positive latitude regions that receive more illumination during northern hemisphere summer on the comet. A broad region of high CO₂/H₂O ratio occurs at negative latitudes in the winter hemisphere, likely the result of deep minima in the H₂O signal (such as the one shown in Fig. 3 on 18 September at 4 hours). This winter hemisphere of the comet is poorly illuminated by the Sun. With limited illumination, this region of the comet nucleus may be considerably colder than other regions, including the neck and smaller lobe. The temperature at and below the surface of the nucleus may be sufficient to sublimate CO and CO₂ but not sufficient to sublimate water. The weak, periodic illumination of this region may be sufficient to drive CO and CO₂ sublimation, producing the separate CO and CO₂ peak (Fig. 3 at 18 hours). However, the compositional asymmetry in the two H₂O peaks cannot be explained in a similar way and might be the strongest indication for heterogeneity in the comet nucleus. The strong heterogeneity in the coma of comet 67P is likely driven by seasonal effects on the comet nucleus. However, the smaller variation of CO and CO₂ compared with H₂O might indicate that CO and CO₂ ices sublimate from a greater depth, whereas H₂O ice sublimates closer to the surface and experiences more direct temperature differences due to sunlight. Furthermore, that lack of overall correlation between H₂O, CO, and CO₂ implies that the outgassing from the nucleus is not correlated or that CO and CO₂ are not strictly embedded in H₂O. For Tempel 1, material was found in layers and supports the above idea (15).




Fig. 4. The nadir point for each pair of CO₂/H₂O measurements over the time period from 17 August through 22 September was mapped to the model surface. The mapping is shown for the bottom side of the larger of the two lobes of the comet, and cometary latitudes run approximately vertically in this. The layering is due to spacecraft rastering above the comet nucleus. A high ratio is measured for the lower part that is poorly sunlit in southern hemisphere winter. [Shape model credit: ESA/Rosetta/MPS for OSIRIS Team MPS/UPD/LAM/IAA/SSO/INTA/UPM/DASP/IDA].

In summary, the coma composition has been measured over many rotational periods of the comet and a wide range of latitudes with high time resolution and compositional detail. Concentrations of the three molecules change over the rotational period of the comet and indicate a strongly heterogeneous coma. For the most part, H₂O dominates, but CO and CO₂ can at times dominate in the coma. These observations also indicate that there are substantial diurnal and latitudinal variations in the coma. Peaks in the H₂O signal are observed, along with deep minima at high negative latitudes when the neck region of the nucleus is blocked from view of the spacecraft. A separate peak in CO₂ signal occurs when the winter hemisphere of the larger lobe of the comet faces the spacecraft. The diurnal and latitudinal variations suggest that compositional differences in the coma may be seasonal and may indicate different subsurface temperatures in the nucleus.

Further observations may distinguish seasonal effect from nucleus heterogeneity. As the comet approaches the Sun, the overall temperature of the nucleus will increase, and as the seasons change, there may be considerable changes in the H₂O, CO, and CO₂ outgassing, with the current high CO₂/H₂O ratio region shown in Fig. 4. In addition, differences in the sublimation of species similar in sublimation temperatures could demonstrate the extend of heterogeneity in the nucleus independent of seasonal changes.

REFERENCES AND NOTES

1. M. F. A'Hearn, R. C. Millis, D. O. Schleicher, D. J. Osip, R. V. Birch, The ensemble properties of comets: Results from narrowband photometry of 85 comets, 1976-1992. *Icarus* **118**, 223–270 (1995). doi: [10.1006/icar.1995.1190](https://doi.org/10.1006/icar.1995.1190)
2. M. J. Mumma, S. B. Charnley, The chemical composition of comets: Emerging taxonomies and natal heritage. *Annu. Rev. Astron. Astrophys.* **49**, 471–524 (2011). doi: [10.1146/annurev-astro-081309-130811](https://doi.org/10.1146/annurev-astro-081309-130811)
3. N. Dello Russo *et al.*, Compositional homogeneity in the fragmented comet 73P/Schwassmann-Wachmann 3. *Nature* **448**, 172–175 (2007). doi: [10.1038/nature05908](https://doi.org/10.1038/nature05908); pmid: [17625560](https://pubmed.ncbi.nlm.nih.gov/17625560/)
4. H. U. Keller *et al.*, First Halley Multicolour Camera imaging results from Giotto. *Nature* **321** (6067s), 320–326 (1986). doi: [10.1038/321320a0](https://doi.org/10.1038/321320a0)
5. H. A. Weaver, M. J. Mumma, H. P. Larson, D. S. Davis, Postperihelion observations of water in comet Halley. *Nature* **324**, 441–444 (1986). doi: [10.1038/324441a0](https://doi.org/10.1038/324441a0)
6. L. M. Feaga, M. F. A'Hearn, J. M. Sunshine, O. Groussin, T. L. Farnham, Asymmetris in the distribution of H₂O and CO₂ in the inner coma of Comet 9P/Tempel 1 as observed by Deep Impact. *Icarus* **190**, 345–356 (2007). doi: [10.1016/j.icarus.2007.04.009](https://doi.org/10.1016/j.icarus.2007.04.009)
7. M. F. A'Hearn *et al.*, EPOXI at comet Hartley 2. *Science* **332**, 1396–1400 (2011). doi: [10.1126/science.1204054](https://doi.org/10.1126/science.1204054); pmid: [21680835](https://pubmed.ncbi.nlm.nih.gov/21680835/)
8. B. P. Bonev *et al.*, The peculiar volatile composition of comet 8P/Tuttle: A contact binary of chemically distinct cometesimals? *Astrophys. J.* **680**, L61–L64 (2008). doi: [10.1086/589649](https://doi.org/10.1086/589649)
9. M. F. A'Hearn, Whence comets? *Science* **314**, 1708–1709 (2006). doi: [10.1126/science.1137083](https://doi.org/10.1126/science.1137083); pmid: [17170287](https://pubmed.ncbi.nlm.nih.gov/17170287/)
10. K.-H. Glassmeier, H. Boehnhardt, D. Koschny, E. Kürt, I. Richter, The Rosetta Mission: Flying towards the origin of the solar system. *Space Sci. Rev.* **128**, 745–801 (2007). doi: [10.1007/s11214-006-9140-8](https://doi.org/10.1007/s11214-006-9140-8)
11. N. Thomas *et al.*, The morphological diversity of comet 67P/Churyumov-Gerasimenko. *Science* **347**, aaa0440 (2014). doi: [10.1126/science.aaa0440](https://doi.org/10.1126/science.aaa0440)
12. H. Balsiger *et al.*, ROSINA-ROSETTA-orbiter-spectrometer-for-ion-and-neutral-analysis. *Space Sci. Rev.* **128**, 1–21 (2007). doi: [10.1007/s11214-006-8335-3](https://doi.org/10.1007/s11214-006-8335-3)
13. S. Mottola *et al.*, The rotation state of 67P/Churyumov-Gerasimenko from approach observations with the OSIRIS cameras on Rosetta. *Astron. Astrophys.* **569**, L2 (2014). doi: [10.1051/0004-6361/201424590](https://doi.org/10.1051/0004-6361/201424590)
14. B. Schläppi *et al.*, Influence of spacecraft outgassing on the exploration of tenuous atmospheres with in situ mass spectrometry. *J. Geophys. Res.* **115**, A12313 (2010). doi: [10.1029/2010JA015734](https://doi.org/10.1029/2010JA015734)
15. M. F. A'Hearn *et al.*, Deep Impact: Excavating comet Tempel 1. *Science* **310**, 258–264 (2005). doi: [10.1126/science.1118923](https://doi.org/10.1126/science.1118923); pmid: [16150978](https://pubmed.ncbi.nlm.nih.gov/16150978/)

ACKNOWLEDGMENTS

The authors thank the following institutions and agencies, which supported this work: Work at the University of Bern was funded by the State of Bern, the Swiss National Science Foundation, and the European Space Agency (ESA) PRODEX Program. Work at the Max-Planck Institut für Sonnensystemforschung (MPS) was funded by the Max-Planck-Gesellschaft (MPG) and Bundesministerium für Wirtschaft und Energie (BMWi) under contract 50QP1302. Work at Southwest Research institute (SwRI) was supported by subcontract 1496541 from the Jet Propulsion Laboratory (JPL) and under NASA prime contract NNX148F71G. Work at BIRA-IASB was supported by the Belgian Science Policy Office via PRODEX/ROSINA PEA 90020. Work at Imperial College London has been partially funded by the Science and Technology Facilities Council (STFC). This work has been carried out thanks to the support of the A*MIDEX project (ANR-11-IDEX-0001-02) funded by the Investissements d'Avenir French Government program, managed by the French National Research Agency (ANR). This work was supported by CNES grants at IRAP, LATMOS, LPC2E, Laboratoire d'Astrophysique de Marseille (LAM) and CRPG and by the European Research Council (grant 267255). A.B.-N. thanks the Ministry of Science and the Israel Space agency. Work at the University of Michigan was funded by NASA under contract JPL-1266313. Work by J.H.W. at SwRI was supported by NASA JPL subcontract NAS703001TONMO710889. ROSINA would not give such outstanding results without the work of the many engineers, technicians, and scientists involved in the mission, in the Rosetta spacecraft, and in the ROSINA instrument over the past 20 years whose contributions are gratefully acknowledged. Rosetta is an ESA mission with contributions from its member states and NASA. We thank the OSIRIS team for giving permission to use the shape model. We acknowledge herewith the work of the ESA Rosetta team. All ROSINA data are available on request until it is released to the PSA archive of ESA and to the PDS archive of NASA.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/347/6220/aaa0276/suppl/DC1
Materials and Methods
Supplementary Text
References

10 October 2014; accepted 31 December 2014
[10.1126/science.aaa0276](https://doi.org/10.1126/science.aaa0276)

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of July 7, 2015):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/347/6220/aaa0276.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2015/01/21/347.6220.aaa0276.DC1.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/347/6220/aaa0276.full.html#related>

This article **cites 15 articles**, 3 of which can be accessed free:

<http://www.sciencemag.org/content/347/6220/aaa0276.full.html#ref-list-1>

This article has been **cited by** 4 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/content/347/6220/aaa0276.full.html#related-urls>

This article appears in the following **subject collections**:

Astronomy

<http://www.sciencemag.org/cgi/collection/astronomy>

Planetary Science

http://www.sciencemag.org/cgi/collection/planet_sci