

A Review of General Physical and Chemical Processes Related to Plasma Sources and Losses for Solar System Magnetospheres

K. Seki¹ · A. Nagy² · C.M. Jackman³ · F. Crary⁴ · D. Fontaine⁵ · P. Zarka⁶ · P. Wurz⁷ · A. Milillo⁸ · J.A. Slavin² · D.C. Delcourt⁵ · M. Wiltberger⁹ · R. Ilie² · X. Jia² · S.A. Ledvina¹⁰ · M.W. Liemohn² · R.W. Schunk¹¹

Received: 5 March 2015 / Accepted: 28 May 2015 © Springer Science+Business Media Dordrecht 2015

Abstract The aim of this paper is to provide a review of general processes related to plasma sources, their transport, energization, and losses in the planetary magnetospheres. We provide background information as well as the most up-to-date knowledge of the comparative studies of planetary magnetospheres, with a focus on the plasma supply to each region of the magnetospheres. This review also includes the basic equations and modeling methods commonly used to simulate the plasma sources of the planetary magnetospheres. In this paper, we will describe basic and common processes related to plasma supply to each region of the planetary magnetospheres in our solar system. First, we will describe source processes in Sect. 1. Then the transport and energization processes to supply those source plasmas to various regions of the magnetosphere are described in Sect. 2. Loss processes are also important to understand the plasma population in the magnetosphere and Sect. 3 is dedicated to the explanation of the loss processes. In Sect. 4, we also briefly summarize the basic

K. Seki seki@stelab.nagoya-u.ac.jp

- ² Department of Atmospheric, Oceanic and Space Sciences, University of Michigan, Ann Arbor, MI 48109, USA
- ³ Department of Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, UK
- ⁴ Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303, USA
- ⁵ Laboratoire de Physique des Plasmas (CNRS-EP-UPMC-UPSud), Ecole Polytechnique, 91128 Palaiseau Cedex, France
- ⁶ LESIA, Observatoire de Paris, CNRS, UPMC, Université Paris-Diderot, 5 place Jules Janssen, 92195 Meudon, France
- ⁷ Physics Institute, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland
- ⁸ Institute of Space Astrophysics and Planetology, INAF, Rome, Italy
- ⁹ High Altitude Observatory, National Center for Atmospheric Research, Boulder, CO 80212, USA
- ¹⁰ Space Sciences Lab., University of California, Berkeley, CA 94720, USA
- ¹¹ Center for Atmospheric and Space Sciences, Utah State University, 4405 Old Main Hill, Logan, UT 84322, USA

¹ Solar-Terrestrial Environment Laboratory, Nagoya University, Nagoya, Aichi 464-8601, Japan

equations and modeling methods with a focus on plasma supply processes for planetary magnetospheres.

Keywords Planetary magnetosphere · Plasma sources · General processes

1 Sources

There are three possible sources of plasma for a planetary magnetosphere. The first one is the surface of solid bodies (planet and/or its satellites), the second one is the planetary (or satellite in the unique case of Titan) atmosphere/ionosphere, if any, and the third source is the plasma from the solar atmosphere, i.e., the solar wind. In this section, we will review processes related to each the sources, i.e., surface in Sect. 1.1, ionosphere in Sect. 1.2, and solar wind in Sect. 1.3.

1.1 Surface

In this subsection, we will review source and loss processes related to the planetary or satellite surface as illustrated in Fig. 1. Topics to be explained include ion-induced sputtering, chemical sputtering, photon stimulated desorption, micro-meteoroid impact vaporization, adsorption by surface, ion-sputtering and radiolysis in icy surfaces, sputter yields from water ice, binding energies, sticking and bouncing, and energy distributions.

1.1.1 Ion-Induced Sputtering

The impact of energetic ions or neutrals (typically of keV/nucleon energies) onto a solid surface causes the removal of atoms, ions and molecules from the top surface. This process is referred to in the literature as sputtering, in particular nuclear sputtering when nuclear interaction between the impacting ion and the surface atoms cause the particle release, or electronic sputtering when the electronic interaction results in particle release, as discussed below for icy surfaces. Sputtering is a well-studied phenomenon in material science (e.g. Behrisch and Eckstein 2007).

The energy distribution for particles sputtered from a solid, $f(E_e)$, with the energy E_e of the sputtered particle, has originally been given by Sigmund (1969) and adapted for planetary science (Wurz and Lammer 2003; Wurz et al. 2007)

$$f(E_e) = \frac{6E_b}{3 - 8\sqrt{E_b/E_c}} \frac{E_e}{(E_e + E_b)^3} \left\{ 1 - \sqrt{\frac{E_e + E_b}{E_c}} \right\}$$
(1)

where E_b is the surface binding energy of the sputtered particle, typically in the eV range, and E_c is the cut-off energy. The cut-off energy E_c , which is the maximum energy that can be imparted to a sputtered particle by a projectile particle with energy E_i , is given by the limit imposed by a binary collision between a projectile atom or ion, with mass m_1 and the target atom, with mass m_2 (to be sputtered) as

$$E_c = E_i \frac{4m_1m_2}{(m_1 + m_2)^2} \tag{2}$$

An example of energy distributions based on Eq. (1) is shown in Fig. 2.



Fig. 1 A schematic illustration of the surface sources and sinks for the exosphere (from Killen and Ip 1999)





The energy imparted to the sputtered atoms and molecules is significant with respect to typical escape energies from planetary objects and a considerable fraction of the sputtered particles do not return to the planetary surface (Wurz et al. 2007, 2010).

The polar angle distribution of sputtered atoms is generally described by a $\cos^k(\varphi_e)$ law (Hofer 1991), where the exponent k, is usually between 1 and 2, and depends on the structure of the surface and φ_e is the ejection angle relative to the normal. For the rough surfaces typically encountered in planetary application k = 1 is usually chosen (Cassidy and Johnson 2005; Wurz et al. 2007).

The sputter yield is the average number of atoms or molecules removed from the solid surface per incident particle. Ion sputtering releases all species from the surface into space reproducing more or less the local surface composition on an atomic level. Preferential sputtering of the different elements of a compound will lead to a surface enrichment of those elements with low sputtering yields in the top-most atomic layers. However, the steady-state composition of the flux of sputtered atoms will reflect the average bulk composition. Thus, particle sputtering, when operative, will give us compositional information about the refractory elements of the bulk surface.

Sputter yields for the different species can be obtained using the TRIM.SP simulation software (Biersack and Eckstein 1984; Ziegler et al. 1984; Ziegler 2004); see also the recent review on computer simulation of sputtering by Eckstein and Urbassek (2007). TRIM, like many other simulation programmes for sputtering, assumes that the collisions between atoms can be approximated by elastic binary collisions described by an interaction potential. The energy loss to electrons is handled separately as an inelastic energy loss. For typical rock (regolith) surface compositions, the total sputter yield, i.e., all species sputtered from the surface taken together, is about 0.12 atoms per incoming solar wind ion at 400 km s⁻¹, considering protons and alpha particles only (Wurz et al. 2007). This sputter yield is the integral over all emission angles and all energies of sputtered particles. The 5 % alpha particles in the solar wind contribute about 30 % to the sputter yield. Heavier ions in the solar wind (Wurz et al. 2007). CMEs can cause increased sputtering of surface material, because their ion density can be much larger than that of the regular solar wind. In addition, alpha particles are often more abundant in the CME plasma, which increases the sputter yield even more.

1.1.2 Chemical Sputtering

When a surface is bombarded with chemically reactive species, chemical alterations in the surface material have to be considered. Chemical reactions between the rock (or regolith grain) and the surface and impacting ions may form species, which are more loosely bound to the surface and thus more easily sputtered. This causes an increase of the sputtering yield or allows for some other release from the surface. This process is usually referred to as chemical sputtering in the literature. In the context of planetary science Potter (1995) considered chemical sputtering for the first time to explain the Na exosphere on Mercury. When a solar wind proton hits the mineral surface processes like the following

$$2H + Na_2SiO_3 \rightarrow 2Na + SiO_2 + H_2O$$
(3)

may occur that liberate the Na from the mineral compound. If this happens on the surface, or the liberated Na migrates to the surface, the liberated Na can be released from the surface also by thermal desorption or photon stimulated desorption. This process was successfully implemented in a 3D model to explain the three dimensional structure of the Na exosphere of Mercury with very good agreement with observations for the spatial distribution and the density (Mura et al. 2009).

1.1.3 Photon Stimulated Desorption

Photon-stimulated desorption (PSD), sometimes also referred to as photon sputtering, is the removal of an atom or molecule by an ultraviolet photon absorbed at the surface, via an electronic excitation process at the surface. PSD is highly species selective, and works efficiently for the release of Na and K from mineral surfaces. Also water molecules are removed from water ice very efficiently via PSD. PSD is considered the major contributor for the Na and K exospheres of Mercury and the Moon (Killen et al. 2007; Wurz et al. 2010). Since PSD releases only Na and K from the mineral matrix it is not very important for the overall erosion of the surface since it will cease once the surface is void of Na and K. The situation is different for PSD of water for an icy object, where the PSD process can remove the major surface species.

The flux of material removed by PSD, Φ_i^{PSD} , of a species *i* from the surface can be calculated by the convolution of the solar UV photon flux spectrum, $\Phi_{ph}(\lambda)$, with the wavelength-dependent PSD-cross section, $Q_i(\lambda)$,

$$\Phi_i^{\text{PSD}} = f_i N_s \int \Phi_{ph}(\lambda) Q_i(\lambda) d\lambda$$
(4)

where N_s is the surface atom density, and f_i is the species fraction on the grain surface. Equation (4) can be approximated as

$$\Phi_i^{\text{PSD}} = \frac{1}{4} f_i N_s \Phi_{ph} Q_i \tag{5}$$

where the factor 1/4 gives the surface-averaged value. The experimentally determined PSDcross section for Na is $Q_{\text{Na}} = (1-3) \cdot 10^{-20} \text{ cm}^2$ in the wavelength range of 400–250 nm (Yakshinskiy and Madey 1999) and for *K* the PSD-cross section is $Q_K = (0.19-1.4) \cdot 10^{-20} \text{ cm}^2$ in the same wavelength range (Yakshinskiy and Madey 2001). Equation (4) can also be written in terms of the PSD yield, Y_i^{PSD} , per incoming photon

$$\Phi_i^{\text{PSD}} = \frac{1}{4} f_i \Phi_{ph} Y_i^{\text{PSD}} \tag{6}$$

which has been determined for water by Westley et al. (1995) in the laboratory. The PSDyield of water is found to be temperature dependent

$$Y_{\rm H_2O}^{\rm PSD} = Y_0 + Y_1 \exp\left(-\frac{E_{\rm PSD}}{k_B T}\right)$$
(7)

with $Y_0 = 0.0035 \pm 0.002$, $Y_1 = 0.13 \pm 0.10$, $E_{PSD} = (29 \pm 6) \cdot 10^{-3}$ eV, and k_B is the Boltzmann constant (Westley et al. 1995). The temperature dependence is very similar to the one for sputtering of ice (see below), which was found later.

1.1.4 Micro-Meteoroid Impact Vaporisation

The impact of micro-meteorites on a planetary surface will volatilise a certain volume of the solid surface, which contributes to the exospheric gas at the impact site. At Mercury, for example, about one to two orders of magnitude more material than the impactor is released because of the high impact speed for meteorites (Cintala 1992).

The ratio of the maximum ejecta velocity to the primary impact velocity decreases with increasing impact speed. The measured temperature in the micro-meteorite produced vapour cloud is in the range of 2500–5000 K. Eichhorn (1978a, 1978b) studied the velocities of impact ejecta during hypervelocity primary impacts and found that the velocity of the ejecta increases with increasing impact velocity and with decreasing ejection angle, with the ejection angle measured with respect to the plane of the target surface. Such ejecta temperatures are significantly higher than typical dayside surface temperatures, but the corresponding characteristic energies are still lower than for particles that result from surface sputtering. In general, the simulated gaseous material from micro-meteorite vaporisation assumes a thermal distribution (e.g. Wurz and Lammer 2003), i.e., a Maxwellian-like energy distribution with an average gas temperature of about 4000 K. For a rocky planetary object in the solar wind the contributions to the exosphere from ion sputtering and from micro-meteorite impact are about the same (Wurz et al. 2007, 2010).

Most of the meteorites falling onto a planetary object are very small, see for example Bruno et al. (2007) for the Moon and Müller et al. (2002) for Mercury. Micro-meteorite bombardment can be regarded as a continuous flux of small bodies onto the surface, and thus as a steady contribution to the exosphere. However, occasionally larger meteorites may fall onto a surface causing a much larger release of particles into the exosphere. Such a scenario was studied for Mercury by Mangano et al. (2007). They found that for a meteorite of 0.1 m radius an increase in the exospheric density by a factor 10–100, depending on species, for about an hour over the density from sputtering should be observed.

1.1.5 Adsorption by Surface

Most of the material released from the surface falls back onto it. Depending on the species, the surface, and the surface temperature the particle may stick or may bounce back into the exosphere. Metal atoms, chemical radicals and similar species will stick to the surface because they become chemically bound, i.e., their sticking coefficient is S = 1. For example a sputtered oxygen atom will stick, i.e., will form a chemical bond with the atom it lands on. Similarly, metal atoms will bind chemically to the surface site they land on. Exception are the alkali metals, where Na and K are observed often in exospheres, and which are transiently adsorbed on mineral surfaces. The probability adsorption (sticking) on silicate surface was measured Yakshinskiy and Madey (2005) as function of the surface temperature. For sodium they found $S_{\rm Na}(100 \text{ K}) = 1.0$, $S_{\rm Na}(250 \text{ K}) = 0.5$ and $S_{\rm Na}(500 \text{ K}) = 0.2$, and for potassium they found $S_{\rm K}(100 \text{ K}) = 1.0$ and $S_{\rm K}(500 \text{ K}) = 0.9$. Non-reactive chemical compounds will only stick to the surface when they freeze onto it, which is important mostly for icy moons and planetary objects further out in the solar system. For example O_2 will not freeze onto the surfaces of the icy moons of Jupiter, thus remain in the atmosphere after they have been released from the surface. The same is true for noble gases.

1.1.6 Ion-Sputtering and Radiolysis in the Icy Surface

In the outer solar system it is quite common to encounter icy moons embedded in the planetary magnetosphere, hence, subjected to ion bombardment. The ion impacts onto a water-icy surface can cause sputtering, ionization and excitation of water-ice molecules. Following electronic excitations and ionization water-ice molecules can get dissociated; chemical reactions among the water-dissociation products result in the formation of new molecules (e.g. O₂, H₂, OH and minor species) that are finally ejected from the surface to the moon's exosphere in a two-phase process (e.g., Johnson 1990).

1.1.7 Sputter Yields from Water Ice

These processes have been extensively studied and simulated in laboratory (e.g., Shi et al. 1995; Johnson 1990, 2001; Baragiola et al. 2003). The energy deposited to a solid by the impacting ion, called stopping power, has two components: electronic excitation of molecules predominant at higher energies and momentum transfer collisions (elastic sputtering) predominant at lower energies (Sigmund 1969; Johnson et al. 2009). Famà et al. (2008) obtained through laboratory data fitting the total sputter yields (i.e., number of neutrals released after the surface impact of one ion) for different incident ions at different energies. They discriminated the contributions due to the two components that produce the release of H₂O (direct ion sputtering) and of O₂ and H₂ (electronic sputtering and radiolysis). The total sputter yield *Y* depends on the type, *j*, and energy, E_j , of the impacting ion and the surface temperature, *T*, and it can be written in the following form:

$$Y_{total}^{j}(E_{j},T) = Y_{H_{2}O}^{j}(E_{j}) + Y_{diss}^{j}(E_{j},T)$$
(8)

where $Y_{\text{H}_2\text{O}}^{j}(E_j)$ is the sputtering yield of the H₂O molecules, given by:

$$Y_{\rm H_2O}^{j}(E) = 1/U_o \cdot \left(\frac{3}{4\pi^2 C_0} a S_n^{j}(E) + \eta \left(S_e^{j}(E)\right)^2\right) \cos^{-f}(\vartheta)$$
(9)

where $U_o = 0.45$ eV is the surface sublimation energy, C_0 is the constant of the differential cross section $d\sigma$ for elastic scattering in the binary collision approximation (Sigmund 1969), a is an energy-independent function of the ratio between the mass of the target molecules and of the projectile (Andersen and Bay 1981), S_n^j is the nuclear stopping cross section, S_e^j is the electronic stopping cross section, η is a factor that gives the proportionality between electronic sputtering and $(S_e^j(E))^2/U_o$, θ is the incidence angle, and f is an exponent of the angular dependence of the yield (Famà et al. 2008).

 $Y_{diss}^{j}(E_{j}, T)$ in Eq. (8) is the yield associated to the loss of O₂ and H₂, produced on ice after its irradiation by energetic ions, given by:

$$Y_{diss}^{j}(E,T) = 1/U_{o} \cdot \left(\frac{3}{4\pi^{2}C_{0}}aS_{n}^{j}(E) + \eta \left(S_{e}^{j}(E)\right)^{2}\right) \frac{Y_{1}}{Y_{0}}e^{-\frac{E_{d}}{k_{B}T(lat,\varphi)}}\cos^{-f}(\vartheta)$$
(10)

where Y_1 and Y_0 are fitting parameters obtained by laboratory data elaboration (see Famà et al. 2008). Only this second term is temperature dependent. Laboratory measurements have shown that H₂O molecules dominate the total release yield at lower temperatures (<120 K) and O₂ and H₂ at higher (>120 K) temperatures (Johnson 2001).

Since H_2 is eventually lost from ice stoichiometrically, and since the measurements used by Famà et al. (2008) referred to water-equivalent molecules, the total yield for the O_2 ejection can be expressed as follows:

$$Y_{O_2}^j = \left[m_{H_2O} / (m_{O_2} + 2m_{H_2}) \right] \cdot Y_{diss}^j(E_j, T) = 0.5 \cdot Y_{diss}^j(E_j, T)$$
(11)

where $m_{\rm H_2O}$, $m_{\rm O_2}$ and $m_{\rm H_2}$ are the molecular masses of a water, oxygen and hydrogen, respectively (Plainaki et al. 2015). The total number, N_i , of the released molecules of type *i* depends on the product of the energy spectrum of the ion fluxes impacting the surface with the energy dependent yield:

$$N_i = \int_E \sum_j dF^j / dE_j \cdot Y_i^j dE_j$$
(12)

🖄 Springer





Some example of laboratory measured sputtering yields of water ice (see website http://people.virginia.edu/~rej/h2o.html) as a function of energy and impact ion species are shown in Fig. 3 together with the theoretical functions by Famà et al. (2008) function (blue) and the Johnson et al. (2009) function (red).

The sputter yields obtained by laboratory simulations could be different (lower or higher) in the planetary environments since the aggregation status and the purity of the surface material could be different from the sample prepared in the laboratory. Also important is the radiative history of the ice, in fact, the irradiation enhances the sputter yield (Teolis et al. 2005).

1.1.8 Binding Energies, Sticking and Bouncing

The kinetic energy of a water molecule ejected from the surface is affected mainly by the surface binding energy and secondarily by the energy or mass of the impacting ion (Johnson 1990, 1998). Although the sublimation energy of H₂O is 0.45 eV/molecule, the sputtered particle energy distributions for molecular ices tend to have maxima at lower energies than a collision cascade prediction with surface binding energy equal to the normal sublimation energy (Brown and Johnson 1986; Boring et al. 1984; Brown et al. 1984; Haring et al. 1984). Several explanations for this phenomenon have been proposed; the surface may be strongly disrupted with many atoms or molecules leaving at once without experiencing the same binding energy as a single atom leaving a planar surface (Roosendaal et al. 1982; Reimann et al. 1984). In addition, the surface region may be electronically and collisionally excited and the interatomic or intermolecular forces are lower as a result of that excitation (Reimann et al. 1984). The assumption of an 'effective' binding energy for the H₂O molecules equal to $E_b = 0.054$ eV, which was experimentally obtained in the past (Boring et al. 1984; Haring et al. 1984) seems appropriate.

The H_2O and the O_2 molecules released from the surface are set up to ballistic trajectories until they either return to the surface of the body or they escape. Upon return to the surface,

the H₂O molecules stick, while the O₂ molecules get thermalized and bounce back to continue their ballistic travel until electron-impact (see next section) ionizes them (Plainaki et al. 2012, 2013). The average kinetic energy that the O₂ molecules have after impacting the surface is about k_BT , where k_B is the Boltzmann constant and T is the surface temperature.

1.1.9 Energy Distributions

The emitted O_2 molecules have a complex energy distribution consisting of two components. The distribution of the O_2 molecules that escape the gravity of an icy moon (e.g., Ganymede) is assumed to be described by an empirical function (Johnson et al. 1983; Brown et al. 1984) used also in earlier modelling (Plainaki et al. 2012, 2013; Cassidy et al. 2007; Shematovich et al. 2005):

$$\frac{dF}{dE_e} = a_n E_{\rm O_2} / (E_e + E_{\rm O_2})^2 \tag{13}$$

where $E_{O_2} = 0.015$ eV (Shematovich et al. 2005), a_n is the normalization factor and E_e is the energy of the ejected O₂ molecules.

The O_2 molecules that have had at least one contact with the surface form a Maxwellian velocity distribution function with a temperature equal to the surface temperature. On the basis of the above, the overall energy distribution of the exospheric O_2 can be considered to be mainly thermal exhibiting however the high energy tail in Eq. (13) (De Vries et al. 1984).

The energy distribution of the sputtered water molecules is similar to the regolith ion sputtering distribution, given by Sigmund (1969) as discussed in Eq. (1). The major difference to sputtering of rock is that the 'effective' binding energy, E_b , is equal to 0.054 eV (Johnson et al. 2002). The binding energy E_b influences significantly the energy spectrum at low energies, while the high energy tail of the distribution is affected mainly by $E_c(E_i)$ (see Eq. (2)).

Finally, since the energetic and heavy ions of giant planets' magnetospheres can produce a release of up to 1000 water molecules per impacting ion after the interaction with the icy moon surfaces (see Fig. 3), the ion sputtering process of water is often a major contributor to the exosphere population for the outer solar system moons, where surface temperatures are generally around 80–150 K and the solar illumination is low. The spatial distributions of the exospheres are expected to depend mainly on the illumination of the moon's surface, which determines the moon's surface temperature responsible for the efficiency of radiolysis (Famà et al. 2008). At these low temperatures, in fact, the averaged expected contribution of sublimated water-ice to the moon's exospheric density is important only locally, i.e., at small altitudes above the subsolar point (Smyth and Marconi 2006; Marconi 2007; Plainaki et al. 2010). The high rate of release of particles at relatively high energy, produce a net escape from the moon and high surface erosion rates; for example, the erosion rate of the icy moons embedded in the Jupiter magnetospheric plasma radiation is estimated in the range of 0.01–0.1 µm per year (Cooper et al. 2001; Paranicas et al. 2002). Usually, H₂ formed in ice diffuses and escapes much more efficiently than O_2 at the relevant temperatures in the outer solar system, and, in turn, escapes from the icy moons because of their relatively weak gravitational fields (Cassidy et al. 2010). Therefore, the irradiation of icy surfaces can preferentially populate the magnetosphere with hydrogen (Lagg et al. 2003; Mauk et al. 2003), leaving behind an oxygen-rich satellite surface (e.g., Johnson et al. 2009).

1.2 Ionosphere

1.2.1 Ionization Processes

Solar extreme ultraviolet (EUV) radiation and particle, mostly electron, precipitation are the two major sources of energy input and ionization in solar system ionospheres (for details see Schunk and Nagy 2009). Relatively long wavelength photons (>90 nm) generally cause dissociation, while shorter wavelengths cause ionization; the exact distribution of these different outcomes depends on the relevant cross sections and the atmospheric species.

Radiative transfer calculations of the solar EUV energy deposition into the thermospheres are relatively simple, because absorption is the only dominant process. Taking into account the fact that the incoming photon flux and absorption cross sections depend on the wavelength and the different absorbing neutral species have different altitude variations, the decrease in the intensity of the incoming flux after it travels an incremental distance ds_{λ} is:

$$d\mathcal{I}(z,\lambda,\chi) = -\sum_{s} n_{s}(z)\sigma_{s}^{a}(\lambda)\mathcal{I}(z,\lambda)ds_{\lambda}$$
(14)

where $\mathcal{I}(z, \lambda, \chi)$ is the intensity of the solar photon flux at wavelength λ and altitude $z, n_s(z)$ is the number density of the absorbing neutral gas, $s, \sigma_s^a(\lambda)$ is the wavelength dependent absorption cross section of species s and ds_{λ} is the incremental path length in the direction of the flux. Integration of Eq. (14) leads to the following expression for the solar flux as a function of altitude and wavelength:

$$\mathcal{I}(z,\lambda,\chi) = \mathcal{I}_{\infty}(\lambda) \exp\left[-\int_{\infty}^{z} \sum_{s} n_{s}(z) \sigma_{s}^{a}(\lambda) ds_{\lambda}\right]$$
(15)

where, $\mathcal{I}_{\infty}(\lambda)$ is the flux at the top of the atmosphere and the integration is to be carried out along the optical path. The argument of the exponential in Eq. (15) is defined as the optical depth sometimes also called optical thickness, τ , thus:

$$\tau(z,\lambda,\chi) = \int_{\infty}^{z} \sum_{s} n_{s}(z) \sigma_{s}^{a}(\lambda) ds_{\lambda}$$
(16)

and thus Eq. (15) can be written as:

$$\mathcal{I}(z,\lambda,\chi) = \mathcal{I}_{\infty}(\lambda) \exp\left[-\tau(z,\lambda,\chi)\right]$$
(17)

Once the ionizing solar photon flux is known, the photoionization rate for a given ion species $P_s(z, \chi)$ can be written as:

$$P_{s}(z,\chi) = n_{s}(z) \int_{0}^{\lambda_{z}} \mathcal{I}_{\infty}(\lambda) \exp\left[-\tau(z,\lambda,\chi)\right] \sigma_{s}^{i}(\lambda) d\lambda$$
(18)

where λ_{si} is the ionization wavelength threshold and $\sigma_s^i(\lambda)$ is the wavelength dependent ionization cross section for species *s*. Figure 4 shows an example of the production rates calculated for Saturn.



Fig. 4 Calculated production rates for Saturn for a solar zenith angle of 27° . Panel **a** shows the direct phototo-production rates and **b** shows the secondary production rates by the resulting photoelectrons. Note that the electron impact ionization rates are very significant at the lower altitudes (from Kim et al. 2014)

1.2.2 Electron Transport

The transport calculations for electrons in an atmosphere are more difficult than those for EUV radiation because scattering and local sources play an important role. In a collisionless plasma the motion of charged particles in a magnetic field can be considered to consist of a combination of a gyrating motion around the field line and the motion of the instantaneous center of this gyration called the guiding center. When the radius of gyration is small compared to the characteristic dimensions of the field line (the case in many ionospheres), one can just concentrate on the motion of the guiding center. Furthermore in most ionospheric applications steady state condition can be assumed; if one further neglects the presence of external electric fields and the divergence of the magnetic field, the equation for the electron flux Φ simplifies down to:

$$\alpha \frac{\partial \Phi}{\partial x} = \sqrt{\frac{m}{2\varepsilon}} \frac{\delta \Phi}{\delta t} \tag{19}$$

where α is the pitch angle with respect to the magnetic field, *r* the distance along the field line, m_e is the mass of the electron, ε is the energy of the electron and $\frac{\delta \Phi}{\delta t}$ denotes collision induced changes in the flux. This equation is usually solved by dividing the flux into a number of equal angular components or streams. The so called two stream approach is the most commonly used approach and it has been shown, using Monte Carlo calculations (Solomon 1993), that given all the uncertainties associated with differential scattering cross sections, it is generally sufficient to consider only two streams.

Once the electron flux is established, as a function of altitude, the electron impact ionization rate P_s of ion species, s, is given by the following relation:

$$P_s(z) = n_s(Z) \int_{\varepsilon_{sl}}^{\infty} \Phi(z,\varepsilon) \sigma_s^i(\varepsilon) d\varepsilon$$
⁽²⁰⁾

where ε_{sl} is the ionization energy threshold for species *s*.

The transport of superthermal ions and neutral gas particles is even more complicated than that of electrons because additional processes, such as charge exchange and ionization are involved. Recent approaches to obtain 3D values of these ion and/or neutral fluxes have used the so-called direct simulation Monte Carlo (DSMC) method (Bird 1994). This approach is well suited to address this problem and as increasing computing resources become available good, comprehensive and accurate solutions are becoming available. Here again once the ion/neutral fluxes are obtained, the impact ionization rate can be directly calculated using an equation analogous to Eq. (20) above.

1.2.3 Loss Processes and Ion Chemistry

The area of science concerned with the study of chemical reactions is known as chemical kinetics. A chemical reaction in which the phase of the reactants does not change is called a homogeneous reaction and in the solar system upper atmospheres and ionospheres these reactions dominate. Dissociative recombination of O_2^+ with an electron is a typical, so called stoichiometric, reaction:

$$O_2^+ + e \to O + O \tag{21}$$

Reactions that can proceed in both directions are called reversible. Charge exchange between an ion and parent atom and accidentally resonant charge exchange between H and O are such reactions:

$$\mathbf{H}^{+} + \mathbf{H} \rightleftharpoons \mathbf{H} + \mathbf{H}^{+} \tag{22}$$

$$O^+ + H \rightleftharpoons H^+ + O \tag{23}$$

The reactions indicated by Eqs. (21), (22), and (23) are called elementary reactions, because the products are formed directly from the reactants. O⁺, for example can recombine with an electron directly, via radiative recombination, but this process is very slow. In most cases atomic ions recombine via a multi-step process. Two examples of such recombination, via multiple-step processes, are:

$$O^{+} + N_{2} \rightarrow NO^{+} + O$$

$$NO^{+} + e \rightarrow N + O$$
(24)

and

$$H^{+} + H_{2}O \rightarrow H_{2}O^{+} + H_{2}O$$

$$H_{2}O^{+} + H_{2}O \rightarrow H_{3}O^{+} + OH$$

$$H_{3}O^{+} + e \rightarrow H_{2}O + OH$$
(25)

The two-step process indicated in Eq. (24) is important in the terrestrial E-region, and the multi-step one indicated by Eq. (25) is very important in the ionospheres of Jupiter and Saturn.

Given the typical thermospheric and ionospheric temperatures the only chemical reactions likely to occur are the so-called exothermic ones. These are reactions that result in zero or positive energy release. Thus, for example, the reaction of H^+ in the ionospheres of Saturn or Jupiter does not take place with ground state H_2 , because the ionization potential of H_2 is larger than that of H. However, if H_2 is in a vibrational state of 4 or higher, the reaction becomes exothermic and can proceed. This is potentially very important in the ionospheres of Jupiter and Saturn (McElroy 1973; Majeed and McConnell 1996). Similarly, in the terrestrial thermosphere the reaction between ground state N and O_2 is very slow, because of the high activation energy that is needed, but the reaction with the excited atomic nitrogen, in the ²D state is rapid and important. For concrete values for chemical reaction reference data can be found in literature (e.g., Schunk and Nagy 1980, 2009; Nagy et al. 1980; Anicich 1993; Fox and Sung 2001; Terada et al. 2009).

1.2.4 Ionospheric Outflows

When a planet has a global intrinsic field, the ions originating in the ionosphere can escape to space from high-latitude regions such as the cusp/cleft, auroral zone, and polar cap. It is observationally known that ions of ionospheric origin can be one of the most important sources of the plasma in the terrestrial magnetosphere especially in the near-Earth regions (see Chappell 2015 for more details). The outflowing ions along the magnetic field can be categorized into several types of ion outflows, i.e., the polar wind, bulk ion upflow, ion conics, and beams. Detailed reviews of observational aspects and theories of ionospheric outflows can be found in the literature (e.g., Yau and Andre 1997; Andre and Yau 1997; Moore and Horwitz 2007; Chappell 2015). Here we briefly summarize important types of ionospheric ion outflows from a magnetized planet or satellite with atmosphere. A good schematic illustration of these outflows can be found in Fig. 1 of Moore and Horwitz (2007).

Polar Wind The polar wind refers to low-energy ion outflows along the open magnetic field lines in the polar ionosphere, mainly caused by an ambipolar electric field formed by the separation of ions and electrons. To achieve charge neutrality with the lighter and faster upflowing electrons, ambient ions are accelerated by the ambipolar electric field. The polar wind has larger flux in the dayside region, where the outflowing photoelectrons can contribute to the ambipolar electric field. However, the controlling factor of the polar wind outflow rate is still under debate (e.g., Kitamura et al. 2012). A variety of modeling efforts have been made for the polar wind (e.g., Banks and Holzer 1969; Ganguli 1996; Schunk and Sojka 1997; Tam et al. 2007). Observations showed a large flux of O⁺ polar wind, which was not expected by classical theories (e.g., Abe et al. 1996; Yau et al. 2007). Possible additional acceleration mechanisms include the mirror force, pressure gradient, and centrifugal acceleration by plasma convection in the curved magnetic field. The acceleration mechanisms of the polar wind ions can be ubiquitous in the ionospheres of magnetized planets or satellites.

Bulk Ion Upflow The bulk ion upflows refer to the upward ion flow in the low-altitude ionosphere around the F region, which is observed in the auroral zone and cusp (e.g., Ogawa et al. 2008). The bulk ion upflows do not significantly contribute to the outflow flux from the ionosphere, since their energy is usually less than 1 eV and well below the escape energy of heavy ions such as O^+ , O_2^+ , and NO⁺. On the other hand, they are considered important to transport these heavy ions to the high-altitude ionosphere to enable them to undergo additional acceleration in the auroral region and cusp. The mechanisms that cause the bulk ion upflow include the electron heating driven by soft electron precipitation, Joule heating of ions, and frictional ion heating.

Ion Conics Ion conics are named after the typical shape of the velocity distribution function of ion outflows caused by transverse acceleration in terms of the local magnetic field. The transverse ion heating with typical energies from thermal to a few keV are often seen in the cusp region and the auroral zone. The resultant heated ions are called TAIs (transversely accelerated ions). They are often accompanied by electron precipitation, electron density depletions, and a variety of different resonant waves, such as lower hybrid (LH) waves or broadband extremely low frequency (BBELF) waves (Norqvist et al. 1996; Frederick-Frost et al. 2007). Once the ions are heated transversely to the magnetic field, the mirror force can accelerate them further upward by conserving kinetic energy. Thus the resultant ion velocity distribution functions at high altitudes show conical shapes. Various types of ion conics have been observed in the terrestrial ionosphere (e.g., Øieroset et al. 1999). This same process can occur and create ion conics, when there is an energy input, such as electron precipitation, into a planetary ionosphere under an open magnetic field line geometry.

Ion Beam It has been observationally shown that there exist parallel electric fields in the auroral region in both the upward and downward current regions. Their significance for auroral acceleration had been widely discussed (e.g., Mozer et al. 1977; McFadden et al. 1999). The formation of parallel electric field has been also studied theoretically (e.g., Brown et al. 1995; Wu et al. 2002). The static electric potential drop typically up to several kV accelerates electrons downward and cause discrete auroras. The same parallel electric field can accelerate ions upward. The resultant ion outflows become mostly field-aligned energetic beams. It is suggested that a distributed field-aligned potential drop produced self-consistently from a balance between magnetospheric hot ion and electron populations, soft electron precipitations, and transverse heating of ionospheric ions. When the magnetospheric population has significant differential anisotropy between the ion distribution and the electron distribution, significant parallel potential drops can develop (Wu et al. 2002).

1.3 Solar Wind

In addition to the sources detailed above, the solar wind can act as a plasma source for magnetospheres. The character of the solar wind changes significantly with increasing radial distance from the Sun, and this, combined with the contrasting obstacles presented by various planetary magnetospheres, leads to a large variation in solar wind-magnetosphere dynamics and in the degree to which the solar wind can act as a plasma source for a given magnetosphere. The electron density, flow speed, and magnetic field strength in the solar wind near the orbit of the Earth are known to be about 7 cm⁻³, 450 km/s, and 7 nT, respectively. The solar wind mostly consists of protons, but contains about 3–4 % of He²⁺ (Cravens 1997; Gombosi 1998; Schunk and Nagy 2009).

It is well known that interplanetary magnetic field (IMF) lines become increasingly tightly wound with distance from the Sun, as modelled by Parker (1958). The average angle that the interplanetary field lines make with respect to the radial direction increases from ~20° at Mercury's orbital distance of ~0.4 AU (Kabin et al. 2000) through ~45° at Earth (Thomas and Smith 1980), ~80° at Jupiter (Forsyth et al. 1996) to ~83° at Saturn's orbital distance of ~9.5 AU (Jackman et al. 2008). The IMF strength also changes with radial distance, with the strength of the B_R component decreasing approximately as r^{-2} . For example, the IMF at Mercury is much stronger than at Saturn (Burlaga 2001), and this has implications for solar wind-magnetosphere coupling.

The form of interaction between the solar wind and magnetosphere changes with the IMF orientation depending on whether the IMF has a parallel or anti-parallel component to the planetary magnetic field at the subsolar magnetopause. The parallel (anti-parallel) case corresponds to the northward (southward) IMF condition at Earth and vice versa at Jupiter and Saturn where the planetary dipoles are oppositely directed to Earth. In addition to magnetic reconnection between the planetary field and IMF, other important physical processes in terms of the solar wind entry into the magnetosphere include the magnetic reconnection, anomalous diffusion across the magnetopause caused by the Kelvin-Helmholtz instability (KHI), and kinetic Alfvén waves.



Fig. 5 Reconnection geometry for Earth from Gosling et al. (1990). The *left-hand side* shows a southward magnetosheath field and the *right-hand side* the northward magnetospheric field. The current layer is shown as the *shaded boundary* in the center of these two regions. As the fields reconnect (where the two separatrices, S1 and S2 cross) and thread the magnetopause, a region of "open" field allows the entry of magnetosheath particles into the magnetosphere. Additionally, a portion of the population is reflected. Both populations are energized by the process of interacting with the current sheet. On the *right* the magnetospheric counterpart is transmitted through the boundary and a population is again reflected, again both are energized. As the reconnection continues the fields convect away from the site (up and down in this figure), carrying the plasma with them. Owing to the velocity filter effect a layer of electrons is seen further away from the magnetopause on both sides (between E1 and I1 and I2 and E2). Once the ions "catch up," a layer of both electrons and ions is then seen (within I1 and I2). (Caption from McAndrews et al. 2008)

1.3.1 Magnetic Reconnection

The solar wind is thought to enter planetary magnetospheres primarily through magnetic reconnection at the magnetopause (Dungey 1961). Reconnection at the magnetopause accelerates and directs a mixture of magnetosheath and magnetospheric plasma along newly opened magnetic flux tubes down into the cusp (see review by Paschmann et al. 2013). The anti-sunward flow in the magnetosheath carries these open flux tubes downstream where they are assimilated into the lobes of the magnetotail (Caan et al. 1973). Much of the plasma injected down into the cusp mirrors and flows upward into the high latitude magnetotail to form the plasma mantle (Rosenbauer et al. 1975; Pilipp and Morfill 1978). The plasma in this region then $E \times B$ drifts down into the equatorial plasma sheet. In this manner magnetic reconnection between the IMF and planetary magnetic field transfers mass, energy and momentum from the solar wind into the magnetosphere. This dayside reconnection at the Earth (Gosling et al. 1990; McAndrews et al. 2008) is illustrated schematically in Fig. 5.

The rate of magnetopause reconnection is modulated strongly by the magnitude and orientation of the IMF relative to the planetary field and plasma conditions in the magnetosheath adjacent to the magnetopause. More specifically, low-latitude reconnection at Earth's magnetopause is strongly controlled by the magnetic shear angle across the magnetopause with the highest rates being observed for the largest shear angles when the interplanetary magnetic field (IMF) has a strong southward component (Sonnerup 1974; Fuselier and Lewis 2011). This is called the "half-wave rectifier effect" (Burton et al. 1975a). The ultimate reason that reconnection at Earth requires large shear angles, ~ 90 to 270°, is the high average Alfvenic Mach number at 1 AU, i.e., $\sim 6-8$ (Slavin et al. 1984). These high Mach numbers result in a high- β magnetosheath and, generally, thin, weak plasma depletion layers (PDLs) adjacent to the magnetopause (Zwan and Wolf 1976). The typically high- β magnetosheaths at the Earth and the outer planets cause the magnetic fields on either side of the magnetopause to differ largely in magnitude. Under these circumstances, reconnection is only possible for large shear angles, typically larger than 90° (Sonnerup 1974). In contrast, the presence of a strong PDL in the inner magnetosheath naturally leads to magnetic fields of similar magnitude on either side of the magnetopause. For low- β magnetosheaths and well developed PDLs observed at Mercury (Gershman et al. 2013), the near equality of the magnetic field on either side of the magnetopause will allow reconnection to occur for arbitrarily low shear angles (DiBraccio et al. 2013; Slavin et al. 2014) such as observed, for example, across heliospheric current sheets (Gosling et al. 2005) or terrestrial low latitude boundary layer under northward IMF (Phan et al. 2005), where the magnetic fields are also nearly equal on both sides.

At Earth an extensive literature exists describing the empirical relationships between the upstream solar wind and IMF (e.g. Perreault and Akasofu 1978; Bargatze et al. 1985; Burton et al. 1975b). These relationships are all based upon the general formula to calculate the magnetopause reconnection voltage that is:

$$\Phi = v_{sw} B_{perp} L \tag{26}$$

where v_{sw} is the solar wind velocity, B_{perp} is the magnitude of the perpendicular component of the IMF (such that $V_{sw}B_{perp}$ is the motional solar wind electric field), and L is the width of the solar wind channel perpendicular to B_{perp} , in which the IMF can reconnect with closed planetary field lines.

The length, *L*, depends in some way on the properties of the interplanetary medium, and is most frequently taken as some function of the "clock angle" of the IMF. Studies have shown that while dayside reconnection (at Earth) is certainly much weaker for northward than for southward IMF, it does not switch off entirely until the clock angle falls below $\sim 30^{\circ}$ (Sandholt et al. 1998; Grocott et al. 2003). Such empirical functions to quantify the rate of dayside reconnection have in turn been applied at Saturn (Jackman et al. 2004) and Jupiter (Nichols et al. 2006) and integrated over time to estimate the amount of flux opened through reconnection at the dayside.

In recent years, the debate about what determines the reconnection rate at the dayside has intensified, in part due to the wealth of spacecraft data at planets such as Mercury, Jupiter and Saturn, which all represent vastly different parameter spaces and thus are likely to differ from the terrestrial magnetosphere in terms of their level of solar wind-magnetosphere coupling (Slavin et al. 2014). A comprehensive study by Borovsky et al. (2008) for Earth found that the reconnection rate is controlled by four local plasma parameters: B_s (the magnetic field strength in the magnetosphere), ρ_s (the plasma mass density in the magnetosphere), and ρ_m (the plasma mass density in the magnetosphere).

Scurry and Russell (1991) argued that dayside reconnection at the outer planets should have a negligible influence as it would be impeded by the high Mach number regimes there.

This argument was countered by the observations of McAndrews et al. (2008) for Saturn and Grocott et al. (2009) for Earth. Subsequently Lai et al. (2012) interpreted a lack of observation of FTEs at Saturn as lack of reconnection. Most recently, Masters et al. (2012) proposed that the plasma beta conditions adjacent to Saturn's magnetopause can restrict the regions over which reconnection can operate. By way of contrast, reconnection at Mercury's dayside has been found to be much more intense than Earth, is independent of the magnetic field shear angle, and varies inversely with magnetosheath plasma β (DiBraccio et al. 2013). Furthermore, large flux transfer events, relative to Mercury's small magnetosphere, occur at Mercury's magnetopause with typical frequencies of 1 every 8 to 10 s (Slavin et al. 2012b; Imber et al. 2014).

MESSENGER observations at Mercury have found that the rate of magnetic reconnection at the dayside magnetopause is on average three times larger than at Earth (Slavin et al. 2009; DiBraccio et al. 2013). A schematic illustration of Mercury's magnetosphere based on MESSENGER observations can be found in Fig. 1 of Slavin et al. (2009). Further, the rate of reconnection at the magnetopause appears independent of IMF direction with high reconnection rates being measured even for small shear angles (DiBraccio et al. 2013; Slavin et al. 2014). These results at Mercury regarding the relationship between low upstream M_A , plasma- β , magnetic shear angle, and reconnection rate parallel the recent developments regarding PDL formation under low M_A (Farrugia et al. 1995) and reconnection as a function of plasma- β (Phan et al. 2013) at Earth. At Earth the typically high- β magnetosheath limits fast reconnection to IMF orientations that have a southward component, i.e. magnetic shear angles across the magnetopause larger than 90° (i.e. the half-wave rectifier effect). However, during encounters with coronal mass ejections at Earth, the upstream M_A approaches values typical of what is seen at Mercury and similar effects are seen; i.e., lowbeta magnetosheaths and high reconnection rates even for small magnetic shears across the magnetopause (Lavraud et al. 2013).

1.3.2 Kelvin-Helmholtz Instability (KHI)

Another important mechanism of plasma entry from the solar wind to the magnetosphere is anomalous diffusion across the magnetopause at low latitudes, i.e., around the equatorial plane. The solar wind plasma needs to be transported in the direction perpendicular to the local magnetic field to realize the diffusion. It is observationally known that the flank plasma sheet of Earth's magnetosphere becomes colder and denser than usual during prolonged periods of northward IMF (e.g., Terasawa et al. 1997; Borovsky et al. 1998). One mechanism to cause the anomalous diffusion can be represented by the Kelvin-Helmholtz instability (KHI), which is driven by a flow shear between the magnetosheath (shocked solar wind) and the magnetosphere. KHI itself is basically an MHD instability, while the non-linear evolution of KHI vortex can facilitate the cross field diffusion and the mixing of the solar wind and magnetospheric plasmas inside the rolled-up vortex.

A number of mechanisms have been proposed that would cause the plasma mixing inside the vortex. One of the candidate mechanisms is magnetic reconnection inside the vortex triggered by vortex roll-up in the presence of finite in-plane component of the magnetic field (e.g., Nykyri and Otto 2001; Nakamura et al. 2008). Once the magnetosheath and magnetospheric field lines are reconnected, the detached plasma from the solar wind can be transported inside the magnetosphere. Another idea to realize the mixing is turbulent transport of solar wind plasma across the field line for the inhomogeneous density case of KHI (e.g., Matsumoto and Hoshino 2006). When the density gradient between the magnetosheath and magnetosphere sides is large, the secondary instability is excited at the density interface inside the vortex and the laminar flow is changed to turbulence. The secondary instability is



Fig. 6 An example of PIC (particle in cell) simulations of KHI for inhomogeneous density case with the density ratio of 0.1. The initial velocity shear layer was located at Y = 0, whose width was set to λ . Color codes show the mixing rate of magnetosheath particles. The mixing rate is defined so that it is maximized (=1) when the magnetosheath-origin particles from Y > 0 at t = 0 occupy the simulation cell equally with the magnetosheath origin from $Y \le 0$. Snapshots of spatial distribution of the mixing rate at t = 256.6 for electrons and ions are shown in panels (**a**) and (**b**), respectively. Panel (**c**) presents the time evolution of the mixing layer. (Adopted and modified from Matsumoto and Seki 2010)

a kind of Rayleigh-Taylor instability (RTI) where the centrifugal force by the rotation motion inside the vortex acts as a gravitational force in the regular RTI. Development of the secondary instability creates a thin, winding, and elongated interface of the solar wind and magnetospheric plasmas. PIC simulation results show that the turbulent electrostatic fields excited by the secondary RTI facilitate an efficient mixing of collisionless plasmas across the field lines. Figures 6a and 6b show an example of such an elongated mixing interface for electrons and ions, respectively (adopted from Matsumoto and Seki 2010).

These proposed nonlinear theories of KHI provide plausible mechanisms for solar wind transport across the magnetopause. On one hand, a remaining problem has been to explain the cold dense plasma sheet formation with KHI. Another question has been how to form a broad mixing layer of several Earth radii observed at Earth (Wing and Newell 2002), since the proposed mixing is basically limited insider the vortex whose size is expected to be much smaller if one consider a simple KHI vortex without nonlinear vortex paring. Based on large-scale MHD and PIC simulations, Matsumoto and Seki (2010) showed that rapid formation of a broad plasma turbulent layer can be achieved by forward and inverse energy cascades of the KHI. Figure 6 shows an example of the full particle simulations. The forward cascade is triggered by growth of the secondary Rayleigh-Taylor instability excited during the nonlinear evolution of the KHI, while the inverse cascade is accomplished by nonlinear mode couplings between the fastest growing mode of the KHI and other KH unstable modes. As a result of the energy transport by the inverse cascade, the growth rate of the largest vortex allowed in the system reaches a value of 3.7 times greater than that of the linear growth rate and it can create the boundary layer extended over several Earth radii (Fig. 6c).

The KHI is also considered important in Saturn's magnetosphere (e.g., Masters et al. 2009, 2010; Delamere et al. 2013). Given that the corotating flows in the magnetosphere have the opposite (same) directions compared to the shocked solar wind flow in the dawn (dusk) side dayside magnetopause, the occurrence of KHI is expected to be highly asymmetrical, i.e., the dawn side magnetopause has a favorable condition to KHI excitation. Observations of kilometric radiation suggested that the KHI at Saturn's magnetopause tends to



occur in the morning sector (Galopeau et al. 1995). Based on the 3-D MHD simulations, Fukazawa et al. (2007a) show that the KHI vortex is more pronounced for the northward IMF case than the southward case. However, the effects of KHI on the plasma mixing and transport in Saturn's magnetosphere are still far from understood.

2 Transport and Energization of Plasma

There are a number of methods by which plasma can be transported and energized within magnetospheres. We refer the reader to Jackman et al. (2014a) for a comprehensive review of transport and loss processes in the magnetospheres of Mercury, Earth, Jupiter and Saturn. In this section we describe major transport and energization processes which are important to understand how to populate various parts of planetary magnetospheres.

2.1 Axford/Hines Cycle

A key transport mechanism, thought to be at work in slowly-rotating magnetospheres, is the so-called viscous interaction driven model (Axford and Hines 1961; Axford 1964). This involves momentum transfer from the solar wind to the magnetotail via quasi-viscous interaction, particularly at the low-latitude magnetopause. It is illustrated schematically in Fig. 7. This cycle can drive circulation within a closed magnetosphere, provided an appropriate tangential-drag mechanism exists. A major mechanism to enable this interaction is the Kelvin-Helmholtz instability described in Sect. 1.3, driven by flow shear at the magnetopause, which may also be coupled with magnetic reconnection (e.g. Hasegawa et al. 2004; Nykyri et al. 2006).



2.2 Dungey Cycle

A second transport mechanism driven by solar wind interaction is the Dungey cycle. In this cycle, dayside reconnection opens magnetic flux, and the solar wind interaction carries these open magnetic field lines from dayside to nightside, where they are stretched out to form the tail lobes (defined here as the open field line region, while noting that centrifugal confinement of plasma to the equator in rapidly rotating systems can alter this picture somewhat (e.g. Hill and Michel 1976; Ray et al. 2009)). As they are stretched out down-tail, open field lines sink in towards the center plane of the tail, where they reconnect again, closing the flux that was opened on the dayside. The "Dungey cycle timescale" refers to the length of time from the opening of the field lines at the dayside to the closing of the field lines on the nightside. Figure 8 shows the stages involved in the Dungey cycle for the case of Earth, where the timescale is ~ 1 hour (Cowley 1982). The Dungey cycle is also known to operate strongly in the slowly rotating magnetosphere of Mercury, with a timescale of just $\sim 1-2$ minutes (Siscoe and Christopher 1975; Slavin et al. 2012a). The relative importance of the Dungey cycle at the rapidly rotating magnetospheres of Jupiter and Saturn is a topic of some debate. Badman and Cowley (2007) estimated that when active, the Dungey cycle timescale at Jupiter is of order several weeks, whereas at Saturn the timescale is ~ 1 week or more (Jackman et al. 2004). Figure 9 illustrates the combination of the Dungey and viscous-cycle flows in the Earth's ionosphere.

2.3 Rotational Driven Transport and Vasyliunas Cycle

The role of rotation in a planetary magnetosphere may be estimated by considering the superposition of dawn-dusk electric field resulting from the solar wind flow and the radial electric field imposed by the planetary ionosphere (Brice and Ioannidis 1970). The resulting potential is

$$\Phi = -\eta v_{sw} B_{sw} r \sin \varphi - \frac{\Omega B_0 R^3}{r}$$
(27)

where v_{sw} and B_{sw} are the solar wind speed and magnetic field, η the efficiency with which the solar wind field penetrates into the magnetosphere, and B_0 , R and Ω the planetary equatorial magnetic field, radius and rotation rate. This implies that the plasma will $E \times B$ drift along closed paths and in the sense of planetary rotation within a distance

$$r_0 = \sqrt{\frac{\Omega B_0 R^3}{\eta v_{sw} B_{sw}}} \tag{28}$$

For the Earth, this approximation suggests a corotating region inside of 4 R_E , reasonably consistent with the observed size of the Earth's plasmasphere. For Jupiter and Saturn, however, the same calculation suggests a size of over 150 and 50 planetary radii, respectively. This would be larger than the actual size of these planetary magnetospheres. In practice, the observed corotating region occupies most, but not all, of these planetary magnetospheres. Nor are the flows at a rigid corotation speed. At Jupiter they begin to depart from corotation somewhere near the orbit of Europa (10 R_J) (McNutt et al. 1979; Krupp et al. 2001) and at Saturn the flows are 10–20 % of full corotation as close to the planet as 4 R_S (Wilson et al. 2009). An example of application of Eq. (27) to Jupiter's case can be found in Fig. 5 of Delamere and Bagenal (2010).

This corotational flow results in a dramatically different distribution of plasma along magnetic field lines and allows internal plasma sources to drive magnetospheric dynamics. The distribution of plasma along a magnetic field line is determined by the gravitational, centrifugal and ambipolar electric potentials (Siscoe 1977; Bagenal and Sullivan 1981)

$$n_{\alpha} = n_{\alpha,0} \exp\left[-\frac{U(\lambda) + q_{\alpha} \Phi(\lambda)}{kT_{\alpha}}\right]$$

$$U(\lambda) = -\frac{GMm_{\alpha}}{LR\cos^{2}\lambda} + \frac{m_{\alpha}}{2}\Omega^{2}L^{2}R^{2}\cos^{6}\lambda$$
(29)

and the requirement of charge neutrality $\sum q_{\alpha}n_{\alpha} = 0$. The above equations assume a dipole magnetic field and isotropic Maxwellian velocity distributions, but can be appropriately modified to treat any magnetic field geometry, as well as non-Maxwellian distributions (e.g. anisotropic Maxwellians (Huang and Birmingham 1992), kappa distributions (Meyer-Vernet et al. 1995), etc.).

When we consider the electric potential inside the synchronous orbit:

$$\left(\frac{2GM}{3\Omega^2}\right)^{1/3} = \left(\frac{2}{3}\right)^{1/3} R_{sync} \tag{30}$$

where R_{sync} is the radius of synchronous orbit, the potential has a maximum at the equator. Outside this distance, there is a potential minimum at the equator and a local maximum at a latitude:

$$\cos^8 \lambda = \frac{2}{3} \left(\frac{R_{sync}}{LR} \right)^3 \tag{31}$$

As a result, ions produced in the equatorial magnetosphere and inside this "critical distance" will freely precipitate into the planetary atmosphere, while those produced farther from the planet are equatorially trapped. In the case of the Earth, the critical distance would be 5.75 R_E . Since this is outside the corotating plasmasphere, no such equatorial trapping occurs in the Earth's magnetosphere. In contrast, trapping may occur outside 1.96 R_J at Jupiter and 1.62 R_S at Saturn. Thus, the plasma in virtually all of these magnetospheres is equatorially trapped. This "critical distance" has also been identified as a limit for stable orbits of charged dust particles, in the limit $m/q \rightarrow 0$ (Northrop and Hill 1983) and in simulations of ions produced over Saturn's ring plane (Luhmann et al. 2006).

In addition to allowing equatorial trapping, the mid-latitude potential minimum also results in a minimum in electron density. While the exact location of this minimum depends on the ambipolar field, and therefore on the abundance and temperature of the various species, calculations using typical, observed values place it close to the latitude given in Eq. (31). At these latitudes, due to their lower mass, protons are expected to be the most abundant species even though they are not at the equator. An increase in proton abundance with latitude has been observed by the Cassini spacecraft at Saturn (Thomsen et al. 2010), but no clear minimum has been reported, probably due to the very low densities present at these latitudes. At Jupiter, protons represent only a few percent of the equatorial ions and mass-resolved observations are unavailable.

This mid-latitude density minimum and the predominance of protons have strong implications for magnetosphere-ionosphere coupling. The dynamical processes of the lowlatitude magnetosphere are connected to the planetary ionosphere through field-aligned currents. These currents are limited by the availability of charge carriers and are therefore sensitive to the electron density profile along a field line. By finding solutions to a onedimensional Vlasov equation, Ray et al. (2009) showed that the current-voltage relation along a Jovian field line differs significantly from the traditional Knight relation (Knight 1973) (see Eqs. (39) and (40)). The saturation current may be one to two orders of magnitude lower and depends on the conditions at the electron density minimum rather than the equator. Other aspects of magnetosphere-ionosphere coupling are mediated by MHD waves. Wave velocities and propagation times are sensitive to the plasma properties along the field lines. As a result, many aspects of magnetosphere-ionosphere coupling at Jupiter and Saturn depend on the poorly measured mid-latitude plasma.

In the presence of equatorial trapping, any plasma sources in the magnetosphere must be balanced by some loss process. In the case of Jupiter and Saturn, plasma is produced by the ionization of neutrals from satellites (primarily Io and Enceladus), rings and the planetary exospheres. Recombination is not an efficient loss process, and charge exchange does not result in a net removal of ions. The main loss process balancing these plasma sources is centrifugally-driven, radial transport. The corotating plasma experiences an outward, centrifugal force. To first order, this is balanced by magnetic tension. Field lines are stretched under the condition:

$$\frac{1}{\mu_0}(\vec{\nabla} \times \vec{B}) \times \vec{B} = \rho \,\Omega^2 r. \tag{32}$$

This result in a current sheet which resembles that of the Earth's magnetotail in some ways, but which is present at all local times. The stretching of the field lines can be roughly approximated by

$$\frac{B_r}{B_z} \sim \frac{H}{r} \frac{\Omega^2 r^2}{2V_A^2} \tag{33}$$

where V_A is the Alfvén speed, H the thickness of the current sheet and B_r the radial field immediately above or below the sheet.

This balance of centrifugal force and magnetic tension is unstable. The situation is analogous to the magnetized Rayleigh-Taylor instability, where a denser fluid is above a less dense one. In this case, the centrifugal force replaces gravity, and radial transport is driven by a denser plasma inside a less dense plasma (Krupp et al. 2004 and references therein). Time scales for this instability are of order the rotation period of the planet, but may be partially stabilized by considerations such as the Coriolis force and coupling to the ionosphere (Pontius 1997).

In the inner and middle magnetosphere, interchange appears to be the key method by which mass can be transported within magnetospheres. It is a process whereby cool, dense plasma can move outward, to be replaced by hotter, more tenuous plasma moving inward, resulting in a net outward transport of mass. This has been observed both at Jupiter (Thorne et al. 1997; Kivelson et al. 1997; Krupp et al. 2004 and references therein) and Saturn (Hill et al. 2005; Burch et al. 2005). The phenomena are less well-measured at Jupiter, since their typical duration there is shorter and below the 80-s time resolution of the Galileo plasma instrument in almost all cases. Typically, the inward-moving flux tubes are characterized by an abrupt increase in magnetic pressure, the disappearance of thermal plasma, and the presence of a hot, energetic particle population. In the case of older (or more inward transported flux tube) events, flux tubes may contain a mixture of low energy plasma diffusing in and energetic particles curvature-gradient drifting out. Much older events are surrounded by a time-dispersed signature in keV and higher energy particles. This is a result of the superposition of the corotating flow and the particles' curvature-gradient drift (in the direction of corotation for ions and opposite it for electrons). The corresponding outward motion of cold, dense plasma has not been reported.

For the rapidly rotating magnetospheres of the outer planets with their large moon- derived plasma sources, the "planetary wind" or "Vasyliunas cycle" is of critical importance (Hill et al. 1974; Michel and Sturrock 1974; Vasyliunas 1983). This Vasyliunas cycle is driven not by the solar wind, but by the energy transferred to internally generated plasma by the fast rotation of these planets. The plasma created deep inside the magnetosphere is accelerated by magnetic stresses from the ionosphere, gains energy, and moves outward from the planet. Centrifugal forces cause the field lines to stretch. These stretched field lines can form a thin current sheet, across which the closed field lines reconnect. This reconnection simultaneously shortens the field line and (like the Dungey cycle), releases plasma down the tail in the form of a "plasmoid". The stages of this cycle, as viewed in an inertial frame of reference, are illustrated in Fig. 10, the picture originally put forward by Vasyliunas (1983).

2.4 Field-Aligned Potential Drop

Many efforts in theories, simulations and observations showed the role played by magneticfield-aligned electric fields at different locations in the Heliosphere. Significant insights of field-aligned processes, such as particle acceleration, parallel electric fields and currents and their relationships come from numerous observations in the terrestrial magnetosphere at different altitudes along magnetic field lines during the last 50 years. To give examples among others, a few missions that contributed to this field after some of the pioneering spacecraft



Fig. 10 Flow pattern (*left*) and field configuration (*right*) expected for a steady-state planetary wind, first proposed for Jupiter by Vasyliunas (1983)

have flown (see the review by Mozer et al. 1980) are listed hereafter. The long-term US program "Defense Meteorological Satellite Program" (DMSP) maintains satellites orbiting at low altitude (830 km) since 1971. In the decades 1980–2000, the Swedish missions VIKING, FREJA and the NASA mission "Fast Auroral Snapshot Explorer" (FAST) were designed to achieve measurements with excellent time and space resolutions at mid-altitudes (from about 400 to 4000 km altitude). The ESA multi-spacecraft pioneering mission CLUS-TER has been exploring all latitudes and longitudes between typically 4 and 20 Earth radii since 2000 over a time period of more than 15 years. The signatures identified in the terrestrial case provide guidelines to interpret observations in other magnetospheres.

In planetary magnetospheres, where plasmas are collisionless in most regions, the mobility of electrons along magnetic field lines is very high as compared to perpendicular motions mostly driven by large-scale electric fields, magnetic or pressure gradients. Therefore, this high field-aligned mobility contributes to cancel out any potential drop that would appear along magnetic field lines. However, from the mid-70s, observations revealed a secondary peak in the energy spectrum of precipitating electrons in the terrestrial auroral region. Evans (1974) interpreted it as the acceleration by a field-aligned potential difference. Numerous observations have then provided evidence of particle acceleration by parallel electric fields and different processes have been invoked. We first recall that field-aligned particle acceleration does not necessarily imply parallel electric fields, an example being the Fermi acceleration. We then present some of the main classes of processes involving quasi-static and transient parallel electric fields.

2.4.1 Fermi Acceleration

The Lagrangian formulation of mechanics describes the particle motion through "generalized coordinates" and associated "generalized momentum". It allows in particular an easy derivation of the conservation laws for cyclic motions. In magnetized environments, particles are rotating around the magnetic field. The first adiabatic invariant associated to this cyclotron motion is μ :

$$\mu = \frac{1}{2} \frac{m v_{\perp}^2}{B} \tag{34}$$

where *m* is the particle mass and v_{\perp} its velocity in the direction perpendicular to the magnetic field *B*. μ shows that the perpendicular velocity increases with the magnetic field. It is conserved if the magnetic field does not vary in time or evolves slowly relative to the gyration period. At time scales much larger than the cyclotron motion, the particle motion is represented by the guiding center of this cyclotron motion. In an approximately dipolar planetary magnetic field, the magnetic field magnitude increases along magnetic field lines from the apex towards the planet. The conservation of the first adiabatic invariant μ shows that the mirror points are located at the points where the magnetic field is equal to B_m , such that:

$$\frac{1}{B_m} = \frac{(\sin \alpha_0)^2}{B_0}$$
(35)

where α_0 and B_0 are the particle pitch-angle and magnetic field magnitude at a given point along the magnetic field line, for example at the apex. The pitch-angle, α , is the angle between the particle velocity and the magnetic field. The location of the mirror points does not depend on the particle energy but only on its pitch-angle. If the particles do not cross another medium with different properties before reaching their mirror points, they remain trapped in the magnetosphere describing this bouncing motion along magnetic field lines.

The Fermi acceleration along magnetic field lines is related to the second adiabatic invariant. The second adiabatic invariant, also called longitudinal invariant, associated with this bounce motion is I:

$$I = \int_{M_S}^{M_N} p_{\parallel} dl \tag{36}$$

where p_{\parallel} is the particle momentum (mv_{\parallel}) in the direction parallel to the magnetic field, *dl* an elementary distance along the curved magnetic field line, M_N and M_S , the magnetic mirror points in each hemisphere, and the integral is taken along the bounce motion. If the magnetic field does not vary in time or evolves slowly relative to the particle bounce motion, the second adiabatic invariant is conserved. An order of magnitude is given by

$$I \approx m v_{\parallel} L_{SN} \tag{37}$$

where $\langle v_{\parallel} \rangle$ is the average velocity in the direction parallel to the magnetic field and L_{SN} , the total length along the magnetic field line between the two mirror points. If for an external cause, the distance between the two mirror points decreases, the conservation of *I* implies that v_{\parallel} increases: this is the so-called Fermi acceleration along magnetic field lines and it does not involve any parallel electric fields. In planetary magnetosphere, this occurs for example during compression events or substorms. More generally, the Fermi acceleration is considered as an efficient process to explain particle acceleration at shocks or the acceleration of cosmic rays.

2.4.2 Parallel Electric Fields, Currents and Particle Acceleration

While most magnetospheric particles remain bouncing back and forth along magnetic field lines between their mirror points, only particles with mirror points located at ionospheric altitudes or below will reach the ionosphere. Their pitch-angle at the field line apex (see Eq. (33)) will be smaller than a maximum pitch-angle α_c , half-angle of the so-called loss cone:

$$(\sin \alpha_c)^2 = \frac{B_0}{B_I} \tag{38}$$

🖄 Springer

where B_1 and B_0 are the magnetic field magnitude at the ionospheric end and at the apex of the magnetic field line. The loss cone is small: for a dipolar magnetic field decreasing with the cube of the distance, the loss-cone angle is of the order of a few degrees at a distance of 10 planetary radii. In planetary magnetospheres, particles within the loss cone are lost from the magnetosphere due to collisions with the upper atmosphere. These precipitating particles also have the fundamental property to be the only magnetospheric particles capable of carrying field-aligned currents between the magnetosphere and the ionosphere. Conversely, the mirror force is favorable for ionospheric particles. All ionospheric particles that could be extracted from the ionosphere reach the magnetosphere and contribute to carry currents.

Highly conductive magnetic field lines provide an electrodynamic coupling between magnetosphere and ionosphere by connecting both plasmas, by transmitting perpendicular electric fields and by circulating field-aligned currents. Both media, ionosphere and magnetosphere, permanently undergo independent large-scale or local processes that modify their electric field and current distribution at a given time. These modifications are transmitted in the conjugate medium through field-aligned currents where they cause a modification of the electrodynamic parameter distribution, which is transmitted to the conjugate medium trough field-aligned currents current carriers, then the coupling is imperfectly achieved and both media are partially disconnected. In this case, the generation of parallel electric fields represents a way to achieve the required current circulation given that the particle acceleration contributes to the increase in the field-aligned current density to the required value. Such parallel electric fields can be associated with quasi-static structures or with transient processes such as waves.

2.4.3 Quasi-Static Parallel Electric Fields

All developed magnetospheres show evidence of accelerated particles, as for example accelerated electrons precipitating into ionosphere and responsible for auroral light emissions. In the terrestrial magnetosphere, observations show auroral electrons accelerated to keV energies; they move faster than the local Alfvén speed, so that they cannot stay in phase with Alfvén waves. This result led Knight (1973) to consider a simple quasi-static model for field-aligned currents carried by ionospheric and magnetospheric electrons accelerated by a quasi-steady parallel electric potential. From the conservation of the energy and of the first adiabatic invariant, he derived a general current-voltage relationship. For applications to auroral magnetic field lines, where:

$$\frac{e\Delta V}{kT_I} \gg 1$$
 and $\frac{e\Delta V}{kT_0} \ll \frac{B_I}{B_0}$

it simplifies to:

$$j_{\parallel} \sim -en_0 \sqrt{\frac{kT_0}{2\pi m_0}} \left(1 + \frac{e\Delta V}{kT_0}\right) \tag{39}$$

and, if $\frac{e\Delta V}{kT_0} \gg 1$, it becomes:

$$j_{\parallel} \sim -en_0 \sqrt{\frac{kT_0}{2\pi m_0}} \left(\frac{e\Delta V}{kT_0}\right) \tag{40}$$

Deringer

Fig. 11 A schematic illustration of the upward current region adapted from Carlson et al. (1998)



where k is the Boltzmann's constant, e and m_e the electron mass and charge, n_0 and T_0 respectively the magnetospheric electron density and temperature, T_I is the ionospheric temperature, ΔV is the total potential drop between the ionosphere and the magnetosphere: $\Delta V = E_I - E_0$, B_I and B_0 respectively the ionospheric and magnetospheric magnetic fields.

This relation provides an estimate of the field-aligned current density that the plasma can carry between the ionosphere and the magnetosphere without any parallel electric fields $(\Delta V = 0)$. It also shows that the presence of a potential drop allows increasing this threshold value to much larger current densities if required for other reasons (e.g., current continuity, mismatch between the ionosphere and the magnetosphere). Field-aligned currents associated with a positive potential drop are directed upward, which corresponds to auroral observations. Improvements were presented by Chiu and Schulz (1978), who took into account the motion of the ions in such a potential structure and their contribution to field-aligned currents.

Following similar steps, Lyons (1980) demonstrated that discontinuities with div $E \neq 0$ in the large-scale electric field pattern could generate large-scale regions of field-aligned currents, associated with parallel electric fields and electron acceleration. Such discontinuities are known to exist near magnetospheric boundaries (boundaries of the plasma sheets, boundary layers, etc. in large-scale plasma flow inhomogeneities. A discontinuity with: div E < 0(>0) would account for upward (downward) field-aligned currents. A typical width of such structures would be of the order of 100 km in the terrestrial ionosphere, i.e. about 0.01 Earth radius.

Acceleration structures are observed at smaller scales in the auroral zone. For instance, accelerated electron precipitations are observed with a typical shape of inverted V and with widths about ten times smaller (\sim 0.001 Earth radius in the terrestrial ionosphere) than the preceding effect. Such acceleration structures are interpreted as the acceleration due to a U-shaped field-aligned upward potential structure, as illustrated in Fig. 11 adapted from Carlson et al. (1998). The magnetic field near the planet is highly incompressible, resulting in nearly electrostatic structures. Downgoing field-aligned electrons crossing the middle of the structure will gain an energy corresponding to the total upward potential drop, but only a fraction of it if they cross the sides. This effect produces the well-known inverted-V shape for the acceleration structure observed by spacecraft flying below it. Spacecraft crossing at higher altitudes (near the top of Fig. 11) will detect outflowing ions accelerated at energies corresponding to the potential drop below the spacecraft and thus with the typical inverted V shape for the same reasons. They will also observe large convergent electric fields near

the edges of the structure, as shown in Fig. 11. These electric structures are not detected below the spacecraft, implying the presence of an electrostatic shock associated with parallel electric fields at intermediate altitudes as shown in Fig. 11 (see a review by Mozer and Hull 2001). Precipitating electrons and outflowing ions carry upward currents.

Diverging electrostatic shocks are also observed and produce the opposite effects with up-going electrons accelerated to somewhat lower energies than the preceding case, and carrying downward currents. More details can be found in a review by Marklund (2009).

De Keyser et al. (2010) proposed a different mechanism to explain the existence of smallscale quasi-static bipolar (convergent or divergent) electric fields. They considered the case of the field-aligned boundary between a dense region of hotter particles and a diluted region of colder particles, as for example the boundary between the plasmasheet and the lobes. This boundary is approximated as a tangential discontinuity which has a finite thickness of the order of the largest Larmor radius, i.e. that of the hotter ions. The transition width differs for each species and is related to their Larmor radius. The difference between the Larmor radii of the hot ions and the hot electrons will produce a charge separation and thus a polarization electric field perpendicular to the interface. The same occurs for the cold ions and electrons, but their Larmor radii are much shorter. In the absence of any potential structure across the interface, this polarization electric field displays a wider region (related to the hot ion Larmor radius) of smaller magnitude and a smaller region (related to the hot electron Larmor radius) of larger magnitude directed in the opposite direction, so that the integrated electric field over the interface cancels out. This produces the bipolar electric field structure. The presence of a potential across the interface attracts or repels ions and electrons depending on its sign, which in both cases results in a monopolar electric field structure, also observed. The mapping in the ionosphere of this magnetospheric electric field distribution and the closure of the currents in the ionosphere lead to the generation of parallel electric fields and currents.

These quasi-static models are very useful in explaining the observed particle acceleration, field and current signatures related to quasi-static structures. However, they cannot explain observations of transient or highly time-dependent features in the distribution of electric fields and currents.

2.4.4 Transient Acceleration

Accelerated particles and large currents are factors capable of triggering instabilities and of generating waves through wave-particle interactions. These waves contribute to modify in turn the initial particle distribution by energy and pitch-angle scattering of the resonant particles, or by energy and momentum propagation to other regions. As a result, the initial electric currents and fields are modified.

Wave-Particle Interactions and Radiation In ideal MHD, shear Alfvén waves propagate with perpendicular electric fields. They have the property to carry field-aligned currents. When perpendicular scales become too small, the ideal MHD approximation is no longer fulfilled, the waves become dispersive and a parallel electric field appears in so-called kinetic Alfvén waves. In the topside terrestrial ionosphere, parallel electric fields can become very important at altitude below a few Earth's radii (Alfvén resonator). The same is true above Jupiter's ionosphere (Ergun et al. 2006). Numerical simulations suggest that Alfvén waves should evolve towards small scales, with the appearance of a filamentary structure resulting in electrostatic structures such as strong Double Layers (DLs) (Mottez and Génot 2011). High resolution remote sensing of the Io-Jupiter magnetic flux tube based on radio

waves observations have demonstrated the existence of strong DLs (up to ~ 1.5 keV amplitude), which were found to move upwards along the magnetic flux tube at the plasma sound velocity (Hess et al. 2007, 2009).

Paschmann et al. (2003) reviewed typical effects at different frequencies occurring in regions of upward and downward currents of the terrestrial auroral zone. Briefly, electron solitary waves or ELF electric field turbulence are found in downward field-aligned region, associated with divergent electric fields and up-going field-aligned electrons. This is the source region of VLF saucers (whistler emissions) and among the first radio emissions observed in the auroral zone. Large-amplitude ion cyclotron waves and electric field turbulence are found in upward current regions, associated with convergent electric fields and precipitating "inverted-V" events. This is also the source region of auroral radiation, powerful emissions observed in the auroral zones of magnetized planets.

One of the most powerful emissions is the auroral radiation observed above the auroral zone of the magnetized planets. These emissions are primarily driven by precipitating electrons accelerated to keV energies. The generation mechanism is well identified as the Cyclotron Maser Instability (Wu and Lee 1979) and has been extensively studied (see review by Treumann 2006). In situ observations, especially by Viking and FAST, have shown that the source regions are the acceleration regions described in Fig. 11, which are strongly depleted in cold plasma ($f_{pe}/f_{ce} < 0.1$ to 0.3) due to the parallel electric field structure (Roux et al. 1993). The instability appears to be most efficiently driven by quasi-trapped energetic electrons, i.e. keV electrons with velocity mostly perpendicular to the magnetic field. However, this quasi-trapped electron population lies in a region of velocity space which should be empty in a simple adiabatic theory, thus its presence in the auroral zone was suggested to be due to time-varying (or space-varying) parallel electric fields (Louarn et al. 1990). The above filamented Alfvén waves are good candidates, consistent with the filamentary structure of the depleted sources of auroral radio radiation.

Reconnection Acceleration Magnetic reconnection is a well-known example of transient situations. The simplest concept involves a configuration with a "X-point" in a 2D geometry, where the magnetic field vanishes. More complicated configurations are considered with 3D geometries, with guide field. In the "frozen-in" conditions where $E + V \times B = 0$, all points of a given magnetic field line will remain magnetically connected during their motion at the velocity *V*. The magnetic reconnection implies that the magnetic field line has been modified or broken and the existing connection region reconnected with another one. This leads to a global reconfiguration of the magnetic structure. Reconnection is generally considered as the result of a local departure from the "frozen-in" conditions and involves parallel electric fields. The triggering factors differ on the plasma types, near the Sun or in planetary magnetospheres; it is generally difficult to predict the time and location where they occur. One of the distant signatures, well-identified onboard spacecraft, is again the particle acceleration. It is observed in the perpendicular direction mainly near the central part of the plasmasheet or in the parallel direction along the separatrices (Paschmann 2008).

On the magnetopause, reconnection can be accompanied by the development of vortices due to the Kelvin-Helmholtz instability (KHI). This process is known to occur at Earth (see e.g. Hasegawa et al. 2009), Mercury (Sundberg et al. 2012), and Saturn (Delamere et al. 2013). Parallel acceleration of electrons is caused by K-H waves, as strongly suggested at Saturn by the observation of Cyclotron Maser radio emission from the morningside sector of the magnetosphere (Galopeau et al. 1995).

2.5 Non-Adiabatic Acceleration

It is sometimes said that the motion of charged particles is nonadiabatic when the second adiabatic invariant (viz., the action integral $I \equiv m \int V_{\parallel} ds$ associated with the particle bounce motion; see Eq. (33)) is not conserved. This may be the case for instance during substorm dipolarization of the magnetic field lines that can lead to different particle energization depending upon their bounce phase; hence, the formation of bouncing ion clusters (e.g., Mauk 1986). However, in the most general case, the motion of charged particles is defined as being nonadiabatic when the first adiabatic invariant (i.e., the magnetic moment associated with the particle gyromotion Eq. (34)) is not conserved. This may occur either when the length scale of the field variation is comparable to or smaller than the ion Larmor radius (spatial nonadiabaticity) or when the time scale of the field variation is comparable to or smaller than the ion cyclotron period, i.e., temporal nonadiabaticity (e.g., Northrop 1963). Under such conditions, the guiding center approximation is not appropriate to investigate the motion of charged particles and a description based on the full equation of motion is necessary. In the steady state terrestrial magnetosphere, the guiding center approximation may be used to characterize the transport of charged particles in the lobes where substantial centrifugal acceleration (up to a few tens of eV) due to $E \times B$ convection of the magnetic field lines may be obtained (e.g., Cladis 1986). The guiding center approximation also is appropriate in the nearly dipolar region of the inner magnetosphere. As a matter of fact, in this region of space, the second adiabatic invariant often is conserved as well so that an adiabatic bounce-averaged description may be adopted to explore the dynamics of, e.g., ring current and radiation belt particles (e.g., Fok et al. 2006). As for the third adiabatic invariant associated with the particle azimuthal drift about the planet, it is often violated; hence, prominent radial diffusion of the particles takes place.

2.5.1 Spatial Nonadiabaticity

At large distances in the equatorial magnetotail, the magnetic field significantly varies on the length scale of the particle Larmor radius and a gyro-averaged description such as that of the guiding center cannot be applied. To characterize the particle behavior, Sergeev et al. (1983) introduced a scaling parameter K defined as the minimum field line curvature radiusto-maximum particle Larmor radius ratio. Sergeev et al. (1983) demonstrated that, as K becomes smaller than ~8, deviations from an adiabatic behavior gradually develop as identified by, e.g., the injection of trapped particles into the loss cone. In a subsequent study, Sergeev et al. (1993) identified the latitude in the auroral zone where the parallel flux becomes comparable to the perpendicular one, as the projection at low altitudes of the nonadiabaticity threshold in the magnetotail (for given particle species and energy). This latitudinal boundary that is referred to as "Isotropy Boundary" forms a convenient proxy to remotely probe the distant tail topology from low-altitude measurements, as shown for instance by Newell et al. (1998).

The fact that particles may not perform a regular helical motion and actually behave in a nonadiabatic manner in the distended Earth's magnetotail was already uncovered in the pioneering work of Speiser (1965). In the case of a pure neutral sheet such as the selfconsistent one of Harris (1962) with opposite magnetic field orientations on either side of the midplane, Speiser (1965) showed that particles execute rapid oscillations about the midplane and are subsequently lost into the flanks. In the case of a quasi-neutral sheet with a small magnetic field component normal to the midplane, such as that due to the Earth's dipole field, Speiser (1965) showed that the above oscillations are coupled with a slow rotation of the oscillation plane so that particles may be turned back toward the planet instead of traveling into the flanks. Sonnerup (1971) considered the action integral $I_Z \equiv m \int V_Z dZ$ (see Eq. (36)) to characterize the behavior put forward by Speiser (1965) since particle orbits do have some regularity (although not in an adiabatic sense).

Using Poincaré surfaces of section or, equivalently, phase space mapping upon crossing of the midplane, Chen and Palmadesso (1986) examined the dynamics of charged particles in the magnetotail in a more systematic manner. In this latter study, it was shown that the above Speiser orbits actually form one of three distinct classes of nonadiabatic orbits. That is, in the Speiser regime, particles do not experience significant pitch angle scattering upon crossing the neutral sheet and those originating from regions of strong magnetic field may return to such regions after neutral sheet crossing; hence, their denomination as "transient" particles. In the second class of orbits, particles experience prominent pitch angle scattering upon crossing of the neutral sheet. Accordingly, particles originating from regions of strong magnetic fields may remain temporarily trapped near the midplane, while those trapped near the midplane may escape after crossing of the neutral sheet; hence, their denomination as "quasi-trapped" particles. Finally, a third class of orbits consists of particles that remain trapped near the midplane, an example of them being the ideal case of particles with 90° pitch angle at equator (see Fig. 4 of Chen and Palmadesso 1986). Chen and Palmadesso (1986) showed that the phase space is systematically partitioned according to these three distinct orbit classes and that the Speiser regime becomes predominant for specific values of the (normalized) Hamiltonian.

Following the approach of Sonnerup (1971), Büchner and Zelenyi (1989) developed a comprehensive interpretation framework of the particle dynamical behaviors. The formalism put forward in this latter study relies on a piecewise description of the particle motion, considering that it can be viewed as a succession of $I_Z \equiv m \int V_Z dZ$ conserving sequences (see Eq. (36)). In this interpretation framework, at some point during transport toward the neutral sheet, particles cross a phase space separatrix that delineates two different dynamical regimes (viz., crossing and non-crossing of the midplane). In the course of these separatrix crossings, small quasi-random jumps of the invariant I_Z occur as put forward by Neishtadt (1987) (see Fig. 14 of Büchner and Zelenyi 1989). In this approach, the Speiser regime (also referred to as "transient") corresponds to a negligible net change of I_Z ; hence, its denomination as "quasi-adiabatic". In contrast, in the above quasi-trapped regime (also referred to as "cucumber-like" in Büchner and Zelenyi 1989), particles are subjected to significant net changes of I_Z . To describe these I_Z changes, Büchner and Zelenyi (1989) introduced a parameter κ defined as the square root of the minimum field line curvature radius-to-maximum Larmor radius ratio (see, e.g., Eq. (41) of Büchner and Zelenyi 1989). This latter κ parameter, that is now commonly used to characterize the adiabatic character of the particle motion, is the square root of the K parameter of Sergeev et al. (1983). It is also comparable with the dimensionless Hamiltonian used by Chen and Palmadesso (1986) since one has $2H \equiv \kappa^{-4}$. According to the analysis of Büchner and Zelenyi (1989), the particle motion turns nonadiabatic for $\kappa < 3$ (equivalently, K < 8 in Sergeev et al. 1983), and the above regimes with transient (Speiser) and quasi-trapped behaviors are obtained for $\kappa < 1$ (a κ regime that is also referred to as the current sheet limit). Between $\kappa > 3$ and $\kappa < 1$, there exists an intermediate regime where particles do not oscillate about the midplane (because of Larmor radii smaller than the field reversal length scale) but their motion is chaotic.

Delcourt et al. (1994) further explored this intermediate $1 < \kappa < 3$ regime, considering a centrifugal perturbation of the particle motion near the magnetotail midplane. The interpretation framework developed in this latter study is that the adiabatic (magnetic moment conserving) sequences upon approach and exit of the neutral sheet are separated by a critical cyclotron turn during which an impulsive centrifugal force (due to the enhanced field line elongation) perturbs the cyclotron motion of the particles. This so-called Centrifugal Impulse Model that describes a single (prototypical) crossing of the field reversal leads to a characteristic three-branch pattern of magnetic moment variations, viz., (i) at small pitch angles, large magnetic moment enhancements regardless of the particle gyration phase, (ii) at large pitch angles, negligible magnetic moment changes and (iii) at intermediate pitch angles, either magnetic moment enhancement or damping depending upon gyration phase. As κ decreases from 3 toward 1, this three-branch pattern gradually expands in velocity space, consistently with the results of Sergeev et al. (1983) (see Fig. 1 of Delcourt et al. 1996). Repeated crossings of the field reversal (equivalently, repeated applications of the three-branch pattern of magnetic moment variations) lead to a chaotic behavior with prominent dependence upon initial phase of gyration since magnetic moment enhancement and damping are obtained at small and intermediate pitch angles, respectively.

In this respect, using single-particle simulations in a model magnetic field of the magnetotail, Ashour-Abdalla et al. (1992) suggested that the $\kappa \approx 1$ regime leads to enhanced particle trapping and duskward drift, a feature referred to as the "wall" region. This $\kappa \approx 1$ regime lies in the mid-tail at the transition between the nearly dipolar region where the particle motion is adiabatic ($\kappa > 3$) and the distant tail where one has $\kappa < 1$. It corresponds to the onset (K = 8) of nonadiabaticity examined by Sergeev et al. (1983) and the "wall" feature is thus at odds with the "Isotropy Boundary" interpretation framework discussed above with particle injection into the loss cone and subsequent precipitation. However, the three-branch pattern obtained with the Centrifugal Impulse Model suggests that the two behaviors coexist, the "wall" feature corresponding to large magnetic moment enhancements at (relatively) small pitch angles while the "Isotropy Boundary" follows from damping of the magnetic moment at intermediate pitch angles.

The nonadiabatic features discussed above are of paramount importance for the development of thin current sheets that are essential magnetotail elements at Earth and at other planets. In the terrestrial magnetosphere, in situ observations from GEOTAIL, CLUSTER and THEMIS have revealed a number of magnetic field features in the tail current sheet such as flapping, flattening, tilting, waving, twisting and bifurcation. This current sheet can become very thin (with a thickness comparable to the ion inertial length), yielding a metastable state that can lead to current sheet disruption as observed during the expansion phase of substorms (e.g., Mitchell et al. 1990). The formation of nongyrotropic distribution functions in these nonadiabatic regimes also leads to nonzero off-diagonal terms in the pressure tensor and allows for a current sheet equilibrium that does not require a prominent pressure gradient along the tail axis (e.g., Ashour-Abdalla et al. 1994). As for the predominant Speiser regimes obtained within specific $\kappa < 1$ intervals, they follow from resonance between the fast particle oscillation about the midplane (imposed by the opposite orientations of the magnetic field above and below the midplane) and the slow gyromotion (imposed by the small magnetic field component normal to the midplane). In this Speiser regime, particles are subjected to prominent energization owing to large displacement along the dawn-to-dusk convection electric field. This efficient Speiser acceleration can thus lead to large particle flux within limited intervals at high energies (small κ); hence, the formation of "beamlets" traveling down to low altitudes as reported in CLUSTER observations (see, Keiling et al. 2004).

2.5.2 Temporal Nonadiabaticity

It was mentioned above that during the expansion phase of substorms, the second adiabatic invariant may not be conserved (Mauk 1986). Indeed, the short-lived electric field induced

by dipolarization of the magnetic field lines can lead to significant energization of particles that are located in the equatorial vicinity while those located at low altitudes may remain unaffected. Here, violation of the second adiabatic invariant is due to temporal variations of the magnetic field on the time scale of the particle bounce period. Note that this second adiabatic invariant may be violated because of spatial variations of the magnetic field as well, as is the case for instance near the frontside magnetopause where particles evolve from bouncing about the equatorial plane to bouncing about the field minimum in the outer cusp region (Shabansky 1971; Delcourt and Sauvaud 1999).

Still, temporal variations of the magnetic field can also lead to violation of the first adiabatic invariant, a behavior that is obtained whenever the magnetic field varies significantly on a time scale comparable to the particle gyro-period. In this regard, it was shown by Delcourt et al. (1990) that, during dipolarization of the magnetic field lines, violation of the first adiabatic invariant may be obtained for heavy ions (O⁺) that have cyclotron periods of a minute or so in the terrestrial mid-tail. As a result, while protons with small gyro-periods behave in an adiabatic manner (with respect to the first invariant), O⁺ may experience prominent nonadiabatic energization, in a like manner to spatial nonadiabaticity, where protons and O⁺ ions may exhibit $\kappa > 3$ and $\kappa \le 1$, respectively.

Unlike the energization by the large-scale convection electric field that is constrained by the magnitude of the cross-polar cap potential drop (typically, in the 50 kV–150 kV range) so that ions drifting over a few R_E across the steady state magnetotail can gain at most a few tens of keV, there is no well defined limit for the energization that can be achieved from the induced electric field (Heikkila and Pellinen 1977; Pellinen and Heikkila 1978). Delcourt et al. (1990) actually showed that O⁺ energization up to the 100 keV range is readily obtained during substorm reconfiguration of the magnetic field lines. Since this energization occurs in a nonadiabatic manner and goes together with prominent enhancement of the particle magnetic moment, it radically changes the long-term behavior of the particles that may evolve from an open drift path (i.e., connected to the dayside magnetopause) to injection into the ring current and rapid gradient drift around the planet owing to the large energy gain realized (see Fig. 5 of Delcourt 2002).

At Earth, a variety of in situ measurements suggest that such a mass-to-charge dependent energization is at work during substorm dipolarization. Post-dipolarization spectra obtained for O⁺ can be significantly harder than those of protons (Ipavich et al. 1984; Nosé et al. 2000). Observations of energetic neutral atoms by Mitchell et al. (2003) also reveal repeated injections of energetic (above 100 keV) O⁺ in conjunction with auroral break-ups, while no similar injections are obtained for protons. The (temporally) nonadiabatic heating at work here increases when the inductive electric field increases or if the ions are located further away from the inner dipolar region in the equatorial magnetotail, and it may actually occur in regions where spatial adiabaticity is achieved (viz., $\kappa > 3$). Note also that prominent fluctuations of the magnetic field on short time scales may somewhat alter this description and lead to significant nonadiabatic heating of protons as well, as displayed in the GEOTAIL data analysis of Ono et al. (2009). From a general viewpoint, temporal nonadiabaticity critically depends upon the characteristics of the magnetic field transition and one may expect that Mercury's environment with small temporal scales as compared to those at Earth is characterized by specific nonadiabatic responses.

2.6 Pick-up Acceleration and Mass Loading

Ions produced within a flowing plasma are a significant source of energy and a sink of momentum, as well as being a source of plasma. Although sometimes used more generally, the classical pick-up process occurs when the parent neutrals have a velocity different from the $(\vec{E} \times \vec{B})/B^2$ drift of the local plasma. The new ions are then accelerated by the convection electric field and form a ring-beam distribution in velocity space. This distribution is unstable to the cyclotron maser instability and may result in the generation of electromagnetic ion cyclotron waves.

Neglecting the energy lost to these waves, the ions have an energy, in the plasma frame, of $2mv_{rel}^2$ (where v_{rel} is relative velocity between the source neutrals and local plasma) or four times the ram energy of a background ions of the same mass. In many cases, this can be a significant source of plasma heating. In addition, acceleration by the convection electric field initially causes the newly created ion and electron to move in opposite directions, and their guiding centers become separated by a gyroradius. The resulting "pick-up current" is $\vec{J} = \frac{v_{rel}m}{B}\frac{dn}{dt} = \frac{m}{B^2}\frac{dn}{dt}\vec{E}$. This is often treated as a "pick-up" conductivity (Thomas et al. 2004 and references therein). The pick-up current, flowing across the background magnetic field, also acts to slow, or mass-load the plasma.

In one common case, pick-up acceleration, heating and mass-loading may occur without producing a net source of mass or plasma. If the ions are produced through symmetric charge exchange, $X^+ + X \rightarrow X + X^+$, then the newly ionized particle will be accelerated as any other pick-up ion, producing a ring-beam distribution, heating and mass-loading. However, the reaction will also generate a fast neutral which escapes the system. As a result, there is no net change in the ion density.

3 Losses

In previous sections, we have considered the various sources of plasma and their transport and energization processes to supply magnetospheric plasmas. We next consider the ways in which this material can be lost from the system, to "balance" the mass budget. There are a number of methods by which plasma can be lost from magnetospheres.

3.1 Tail Reconnection and Plasmoids

Magnetic reconnection in a planetary magnetotail is a key mechanism by which magnetic field lines stretch to instability and break, which then allows the release of parcels of mass and plasma called plasmoids, of varying sizes and shapes (Hones 1976, 1977). Observations in the Earth's magnetosphere have shown that plasmoids are typically about 1 to 10 Re in diameter (Ieda et al. 1998; Slavin et al. 2003). Figure 12 shows a schematic of the formation of earthward and tailward-moving plasmoids following reconnection. Figure 13 shows the effect that a large plasmoid has on the surrounding magnetotail at Earth including the magnetic field signatures associated with the traveling compression region in the surrounding lobe regions and the tailward retreat of the near-Earth neutral line (NENL) following ejection (Slavin 1998). In situ observations of tail reconnection include observations of changes in magnetic field topology and plasma flows. In recent years the study of tail reconnection has been extended beyond Earth. Plasmoids have been observed in the magnetotails of Saturn (e.g. Jackman et al. 2007, 2011, 2014b), Jupiter (Russell et al. 1998; Vogt et al. 2010, 2014), and Mercury (Slavin et al. 2009, 2012b; DiBraccio et al. 2013).

In recent years several authors have sought to consider the role of tail reconnection as a loss mechanism for magnetospheric plasma (e.g. Bagenal and Delamere 2011). At Jupiter, Bagenal (2007) highlighted the mismatch between the inferred mass input rate from Io of \sim 500–100 kg/s and the mass loss rate from plasmoids, estimated at \sim 30 kg/s. Kronberg



et al. (2008) attempted a similar calculation (based on Galileo energetic particle measurements) and found that their inferred mass of $\sim 8 \times 10^5$ kg per plasmoid would require far more plasmoids than had been observed to account for the input. Vogt et al. (2014) completed the most comprehensive study to date at Jupiter, whereby they found that mass loss ranged from $\sim 0.7-120$ kg/s. They concluded that while tail reconnection is indeed an active process at Jupiter, it likely cannot account for the mass input from Io, suggesting that additional mass loss mechanisms may be significant. Jackman et al. (2014b) investigated the analogous picture at Saturn. They found an average mass loss rate of ~ 2.59 kg/s, much less than the ~ 100 kg/s expected to be loaded into the magnetosphere by the volcanic moon Enceladus.

These studies raise the question: If large-scale reconnection is not sufficient to account for the required loss of material from the tails of Jupiter and Saturn, what other processes/new physics are required to balance the mass budgets? Other loss mechanisms are investigated in the sections below.

3.2 Charge Exchange

In the Earth's magnetosphere, there exist a region called the ring current, where high energetic ions and electrons with energy between hundreds of eV and hundreds of keV are trapped by Earth's dipole-dominated magnetic field (Frank 1967; Williams 1981). In the ring current, the ions (electrons) drift westward (eastward) due to the magnetic drift, and the ring current development causes the decrease in the horizontal magnetic field component at Earth's surface. Thus, the strength of the ring current is often measured by the Dst or SYM-H indices derived from ground-based magnetometer observations (Sugiura 1964; Wanliss and Showalter 2006). If the magnetic field is strong enough and dominated by the dipole component and there are transportation and energization processes to populate highenergy ions in the inner magnetosphere of a planet, a similar ring current is expected to exist at such planets.

One efficient loss mechanism for the terrestrial ring current particles is the charge exchange (see Eqs. (22) and (23)) of the ring current ions with the neutral hydrogen that makes up the geocorona. When the convection weakens, this becomes the dominant process by which ring current ions are removed from the system, depleting the inner magnetosphere of its energetic population. The geocorona is a halo-like extension of the exosphere out to several Earth radii, consisting of relatively cold (\sim 1000 K), very tenuous neutral hydrogen atoms with densities ranging from thousands of atoms per cubic centimeter at the inner edge of the ring current to less than a hundred at geosynchronous orbit. This cold gas plays a critical role in the energy budget of the Earth's inner magnetosphere since the charge exchange reactions make the exosphere act as an energy sink for ring current particles, replacing a hot ion with a cold one. Singly charged ring current ions can be neutralized after collisions with thermal exospheric hydrogen atoms as described below:

$$\mathrm{H}^{+} + \mathrm{H}_{cold} \to \mathrm{H} + \mathrm{H}_{cold}^{+} \tag{41}$$

$$O^+ + H_{cold} \to O + H^+_{cold} \tag{42}$$

$$\mathrm{He^{+}} + \mathrm{H}_{cold} \to \mathrm{He} + \mathrm{H}_{cold}^{+} \tag{43}$$

The incident ring current ion picks up the orbital electron of the cold geocoronal hydrogen atom resulting in the formation of an Energetic Neutral Atom (ENA). These particles are not affected by magnetic or electric field forces therefore they are no longer trapped in the geomagnetic field and leave the interaction region in ballistic orbits in the direction of the incident ion velocity at the time of the impact. If the resulting ENA's velocity exceeds the Earth's gravitational escape field, then it is lost into space or precipitates down into the ionosphere. On the other hand, the low energy ENAs populate the plasmasphere. Meinel (1951) first reported the existence of energetic neutral atoms, based on observations of precipitating energetic neutral hydrogen precipitating into the upper atmosphere during auroral substorms. A few years later, Dessler and Parker (1959) were the first to suggest that charge exchange between protons and neutral atmospheric hydrogen atoms would effectively contribute to the decay of the ring current, although the effectiveness of ion removal from the ring current through charge exchange processes was previously investigated by Stuart (1959) and Fite et al. (1958).

Multiply charged ions allow for multiple charge exchange reactions,

$$\mathrm{He}^{++} + \mathrm{H}_{cold} \to \mathrm{He}^{+} + \mathrm{H}^{+}_{cold} \tag{44}$$



and Spjeldvik and Fritz (1978) showed that the higher charge states of helium and oxygen ions are increasingly important for energies above 100 keV, while at energies below this cutoff the lower charge states are dominant. Energetic neutral atoms generated in the main ring current traversing the inner magnetosphere can be re-ionized. This happens by converting ENAs back into ring current ions albeit on new L shells, undergoing subsequent charge-exchange collisions with geocoronal atoms and generating secondary ENA fluxes that can participate in further ionizing collisions (Bishop 1996). This yields the formation of a secondary ring current close to the Earth, at L shell values of approximately 3, although this is not a large ring current population. Moreover, low pitch angle ions are subject to additional charge exchange collisions with the oxygen atoms in the upper atmosphere.

Solar far-ultraviolet light is reflected off this hydrogen gas (Chamberlain 1963) and so its abundance has been quantified. It has been reported (Fahr 1974; Rairden et al. 1986; Hodges 1994; Østgaard et al. 2003; Fuselier et al. 2010; Zoennchen et al. 2010, 2011; Bailey and Gruntman 2011) that the geocoronal hydrogen density decreases exponentially with radial distance. This means that at large altitudes down the magnetotail, the collisions with the neutral hydrogen become negligible. However, in the ring current region, these collisions become increasingly important and magnetospheric H^+ can be easily removed by charge exchange with the neutral exospheric hydrogen.

The probability of collisions with neutral atoms from the exosphere depends strongly on the energy of the incident particles and is determined by the charge exchange cross sections. Charge exchange cross sections are both energy and species dependent and thus different ring current ion species have different charge exchange lifetimes. A compilation of charge exchange cross sections for various ring current ions can be found in Spjeldvik (1977), Smith and Bewtra (1978), and Orsini and Milillo (1999).

Numerous studies, both based on both observations and numerical modeling show that due to the strong species and energy dependence of the charge-exchange cross sections along with the temporal and spatial dependence of ring current composition, the charge exchange process strongly affects the ring current plasma. Figure 14 shows the profile of charge exchange lifetime as a function of energy and species (Liemohn and Kozyra 2005). Moreover, it is inferred that the charge exchange loss processes are predominantly important after the initial phase of the ring current decay.

The efficiency of ion removal from the ring current through charge exchange depends on several factors: the energy and the species of the ion population as well as the density of the neutral cloud. The latter depends on the changes in the atmospheric temperature and density, the radiation pressure exerted by the solar far ultra violet photons and the strengths of all these interactions determine the structure of the exosphere. Therefore reliable measurements of the geocoronal density are essential in determining the relative importance of charge exchange losses of ring current ions. The majority of geocoronal models report on vastly different densities in the inner magnetosphere (Ilie et al. 2013) and therefore the decay rates and lifetimes for ring current ions are significantly different depending on the neutral

density distribution, affecting the amount of ENAs emitted in a given region in space (see Fig. 15).

Keika et al. (2003, 2006), based on measurements of energetic neutral atoms (ENAs) made by the High Energy Neutral Atom (HENA) imager on board the Imager for Magnetopause-to-Aurora Global Exploration (IMAGE) satellite, show that the rate of the charge exchange energy losses is comparable to the ring current decay rate for the intervals of the slow decay, while the loss rate is much smaller than the decay rate in the rapid decay phase, in particular for the early stage of a storm recovery. Similarly, Jorgensen et al. (2001) show that during the fast recovery the measured ENAs can only account for a small portion of the total energy loss and the lifetime of the trapped ions is significantly shorter during the fast recovery phase. Furthermore Kozyra et al. (2002) suggested that charge-exchange losses can be solely responsible for the decay of the ring current during the recovery phase only if IMF abruptly turns northward at the end of the main phase.

The neutral gases in the upper atmospheres of Jupiter and Saturn are molecular and atomic hydrogen and thus either as a result of direct ionization or dissociative ionization a significant number of H^+ ions are created. H^+ can only recombine directly via radiative recombination, which is an extremely slow process and thus there must be other ways to remove them otherwise very large ion densities would result. As indicated in Sect. 1.2.3 (Loss Processes and Ion Chemistry), McElroy (1973) suggested some time ago that the following charge exchange would be important in removing H^+ :

$$\mathbf{H}^{+} + \mathbf{H}_{2} \ (v \ge 4) \longrightarrow \mathbf{H}_{2}^{+} + \mathbf{H}$$

$$\tag{45}$$

 H_2^+ is rapidly transformed to H_3^+ via the following reaction:

$$H_2^+ + H_2 \longrightarrow H_3^+ + H \tag{46}$$

 H_3^+ will most likely undergo dissociative recombination and thus this series of reactions removes ions relatively rapidly. There is another way that H^+ can be lost at Jupiter and Saturn (see Eq. (25)), namely by reacting with water molecules, originating in the rings (Connerney and Waite 1984).

3.3 Precipitations into Planets

3.3.1 High Latitudes

As seen in Sect. 2.4, the atmospheric loss cone can be defined at any location by its halfangle sin $\alpha_{lc} = (B/B_m)^{1/2}$ (see Eq. (34)) where *B* is the magnetic field amplitude at the position considered and B_m its value at the ionospheric end of the magnetic field line. Charged particles with pitch angle $< \alpha_{lc}$ will precipitate into the planet and be lost for the magnetosphere. The loss cone is permanently fed by new particles resulting from processes such as pitch-angle scattering by electric and magnetic fluctuations (e.g. whistler waves; see Bolton et al. 2004 and references therein). Due to the converging field line geometry, most precipitations occur at relatively high magnetic latitude ($\sim 55^{\circ}-75^{\circ}$).



Fig. 15 Color contours of lifetimes of H^+ (*left column*) and O^+ (*right column*) as a function of energy and radial distance from the Earth. From *top* to *bottom* are shown the lifetime predictions from Rairden et al. (1986), Østgaard et al. (2003), Hodges (1994), Zoennchen et al. (2011) and Bailey and Gruntman (2011). The color scale is logarithmic and lifetimes are in seconds. Figure from Ilie et al. (2013)

Auroral Ovals Precipitations of electrons with energy ≥ 0.1 keV and of protons or ions with energy $\ge a$ few keV produce auroras (Birkeland 1910), seen from the ground as curtains of light, and from space as bright variable narrow circumpolar ring, arcs and spots. The precipitating electrons have energies in the range $\sim 100 \text{ eV}-10 \text{ keV}$ for the Earth (Feldstein et al. 2001) and Saturn (Cowley et al. 2004), reaching more than 100 keV for Jupiter (Prangé et al. 1998). This is well above their thermal energy in the magnetosphere or solar wind ($\le 1 \text{ eV}$), thus strong acceleration is required, as discussed in Sect. 2.4. Total precipitated auroral power is up to $\sim 10^{11}-10^{12}$ W for the Earth and Saturn, $\sim 10^{13}-10^{14}$ W for Jupiter (Clarke 2012).

X-ray to radio emissions are produced in the high altitude atmosphere (80–300 km on Earth, 10⁻⁵–10⁻⁹ bar at Jupiter) or in the precipitating beam (Prangé 1992; Bhardwaj and Gladstone 2000). The visible aurora is most spectacular on Earth, related to the excitation/deexcitation of O (red and green lines), N (blue line) and N₂ (purple), whereas H- α and H- β lines are very faint at Jupiter. The UV aurora, 10× to 100× more intense than visible ones, result from the collisional excitation (by electrons from a few to 100 keV) and then radiative deexcitation of N_2^+ , N, H at Earth, and H (Ly- α) and H₂ (Lyman and Werner bands) at Jupiter. The X-ray aurora on Earth is mainly generated via bremsstrahlung from precipitating electrons, and at Jupiter from the collisional excitation (followed by radiative deexcitation) of deep internal levels of O and S ions by precipitating heavy ions of energy >100 MeV. The X and UV aurorae are often pulsed on timescales of tens of minutes. The IR auroral emission is due to atmospheric Joule heating (followed by radiative cooling). It is emitted as nitrogen lines at Earth and H₃⁺ and hydrocarbons lines at Jupiter. As UV absorption by hydrocarbons is strongly frequency-dependent, the comparison between auroral and laboratory H and H_2 UV spectra provides information to deduce the depth at which precipitated energy is deposited and, with an atmospheric model, to derive the nature and energy of precipitating particles. Coherent circularly polarized cyclotron radio emissions are generated below $\sim 1 \text{ MHz}$ ($\leq 40 \text{ MHz}$ at Jupiter) by the interaction of unstable precipitating (or mirrored) energetic (1–10 keV) electron populations with electromagnetic fluctuations, in a rarefied and magnetized plasma $(f_{pe}/f_{ce} \ll 1)$ (Zarka 1998). Their generation causes the diffusion of the electrons in velocity space (Pritchett 1986) in particular into the loss cone, causing further precipitations. Imaging the auroral activity in UV (HST-Prangé et al. 1998), IR (ground-based telescopes—Connerney et al. 1993) and radio (via DE-1 (Huff et al. 1988) or Cassini spacecraft (Cecconi et al. 2009)) permits to map the precipitations and, by projection along the magnetic field, the magnetospheric activity.

Polar Cusps and Satellite-Magnetosphere Interactions In addition to the auroral ovals, at the limit between open and closed field lines at Earth or near the corotation breakdown region at Jupiter and Saturn, signatures of precipitations are also observed at the magnetic footprints of the polar cusps and of satellites embedded in the giant planets' magnetospheres (Waite et al. 2001; Pallier and Prangé 2004). Cusp signatures are around 12:00 LT and reveal sporadic dayside reconnections at timescales between 5 min (at Earth) and 20 min (at Jupiter), causing direct precipitation of accelerated particles in the polar cusps. They are more intense for a southern solar wind B_z at Earth (northern at Jupiter). The auroral input power into the cusp is only ≈ 1 % of the total auroral input power. The magnetic footprints of Io, Ganymede and Europa were detected in UV at Jupiter (Bonfond 2012), as well as that of Enceladus at Saturn (Pryor et al. 2011). Precipitation in the satellites magnetic flux tubes result from the imposed current across the satellite due to the electric field $E = v \times B$ arising from the motion of the satellite (at velocity $v = v_{Keplerian} - v_{corotation}$) across the planetary magnetic field lines. This current is carried by Alfvén waves accelerating electrons. In

the Io-Jupiter case, the precipitated power reaches 10^{12} W, i.e., $\sim 10-15$ W/m² at the satellite ionospheric footprints. This power, within a factor 2 of the solar input, strongly heats the local ionosphere and modifies its properties, such as conductivity (Prangé et al. 1996). Satellite footprints have downstream tails related to currents reaccelerating the magnetospheric plasma downstream of the obstacle.

Magnetospheric particles also precipitate onto the surface of embedded satellites. If the latter possesses a magnetic field, precipitating particles are guided toward the magnetic poles of the satellite, generating satellite auroras as well as significant surface alterations, as for example, in the case of Ganymede's polar caps (Khurana et al. 2007).

3.3.2 Low Latitudes

Radiation Belts and Synchrotron Losses Radiation belts consist of electrons and ions accelerated to very high energies (0.1 to > 10 MeV) and brought by radial inward diffusion close to the planet (typically between the surface and ~ 6 radii), where they bounce between their mirror points. Satellites and rings embedded in the belts cause strong collisional absorption of these energetic particles. Unabsorbed electrons can emit synchrotron radiation, a linearly polarized incoherent nonthermal radiation from high energy electrons in cyclotron motion in a magnetic field. This emission extends over a spectral range from <100 MHz to several GHz in the case of Jupiter, and can thus be imaged by ground-based radiotelescopes (Bolton 2004). Intensity is maximal near the equator (trapped population) and near the poles (mirror points, where the residence time is maximum due to low parallel velocity). The lifetime of an emitting electron is relatively short (10^8 to 10^9 s), during which the perpendicular energy of the particle is radiated away and finally causes precipitation onto the planet at low latitudes (≤50°). At Earth and Saturn, synchrotron emission (yet undetected) may exist at much lower frequency and intensity. Saturn radiation belts are largely absent due to ring absorption, but a small belt was discovered by Cassini between the inner edge of the rings and the planet (<1.4 Rs) (Krimigis et al. 2005).

Precipitations from the Rings Other precipitation into Saturn's ionosphere come from the rings' ionized atmosphere (Luhmann et al. 2006). It is composed of O_2^+ and O^+ ions between ~1.4 and ~2.4 *Rs* near the equator, resulting from the ionization by sunlight and magnetospheric impacts of the neutral atmosphere due to sputtering, photo-desorption and meteoroid impacts. The ion motions in the planetary quasi-dipolar magnetic field, subject to the corotation electric field, gravitation and collisional scattering, lead to precipitation into the planet at mid-latitudes (30°–40°) of ions created at radial distances within the corotation orbit at ~1.8 *Rs*. Due to the slight North-South asymmetry of the magnetic field (stronger in the northern hemisphere), precipitation (of energy $\leq 100 \text{ eV}$) occurs mostly in the southern hemisphere.

4 Basic Equations and Modeling Methods

4.1 MHD (Magnetohydrodynamic) Simulation

The basic equations of magnetohydrodynamics (MHD) are derived in numerous textbooks including those by Chen (1984) and Krall and Trivelpiece (1986) and are traditionally presented in terms of the primitive or state variables; density (ρ), velocity (u), thermal pressure

(P), and magnetic field (B) as

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \vec{u}) = 0, \tag{47}$$

$$\rho \frac{\partial \vec{u}}{\partial t} + \vec{u} \cdot \nabla \vec{u} + \nabla P - \frac{1}{\mu_0} \nabla \times \vec{B} \times \vec{B} = 0, \tag{48}$$

$$\frac{\partial P}{\partial t} + \gamma \nabla (P\vec{u}) - (\gamma - 1)\vec{u}\nabla \cdot P = 0, \tag{49}$$

$$\frac{\partial B}{\partial t} - \nabla \times (\vec{u} \times \vec{B}) = 0.$$
(50)

The assumption of ideal gas law has been used to define the pressure equation (49) and the fact that the current density (J) is the curl of the magnetic field has been used to simplify the equations. More importantly in the generalized Ohm's law,

$$\vec{E} = -(\vec{u} \times \vec{B}) + \eta \vec{J} + \frac{1}{en_e} \vec{J} \times \vec{B} - \frac{1}{en_e} \nabla P_e,$$
(51)

terms related to the finite resistivity (η), Hall effect ($J \times B$), and electron pressure (P_e) have been neglected to get to Eq. (50). This formulation is commonly referred to as the equations of ideal MHD and it is important to point out that unless some term in the generalized Ohm's law is restored either analytically or numerically it is not possible for magnetic reconnection to occur in a system that obeys the equations of ideal MHD.

Numerical simulation of these equations usually involves discretization in space and time so it is common to formulate the ideal MHD equations in conservative form in order to allow for the direct application of advanced numerical techniques. The algorithm paper by Toth et al. (2012) not only provides a description of the motivation for utilizing conservative formulation but it also provides a more detailed discussion of the Hall and multifluid formulations than can be covered here. The conservative formulation involves equations of the form,

$$\frac{\partial U}{\partial t} + \nabla \cdot \vec{F}(U) = 0, \tag{52}$$

so that on a discrete grid the change of a conserved quantity is simply the sum of fluxes entering and leaving that cell. Recasting the ideal MHD equations in conservative form results in,

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \vec{u}) = 0, \tag{53}$$

$$\frac{\partial \rho \vec{u}}{\partial t} + \nabla \cdot \left(\rho \vec{u} \vec{u} + \left(P + \frac{B^2}{2\mu_o} \right) - \frac{\vec{B}\vec{B}}{\mu_o} \right) = 0, \tag{54}$$

$$\frac{\partial \mathcal{E}}{\partial t} + \nabla \cdot \left(\vec{u} \left(\mathcal{E} + P + \frac{B^2}{2\mu_o} \right) - \vec{u} \cdot \frac{\vec{B}\vec{B}}{\mu_o} \right) = 0, \tag{55}$$

$$\frac{\partial B}{\partial t} + \nabla \cdot (\vec{u}\vec{B} - \vec{B}\vec{u}) = 0, \qquad (56)$$

🖄 Springer

where

$$\mathcal{E} = \frac{P}{\gamma - 1} + \frac{\rho U^2}{2} + \frac{B^2}{2\mu_o}$$
(57)

is the total energy density of the plasma element. In this formulation it is clear that the change in momentum density in a given region or computational cell is the result of the momentum entering or leaving the cell combined with the effects of thermal and magnetic pressure forces as well as with magnetic tension. Along with these equations comes an important constraint from Maxwell's equations, namely, the fact that the magnetic field must be divergence free ($\nabla \cdot B = 0$) throughout the entire computation domain for all times. In computational solvers this means using a simple projection scheme, a staggered type mesh (Yee 1966) with the magnetic fluxes defined on the faces and the electric fields on the edges, or the constrained transport 8-wave scheme (Powell et al. 1999). The staggered mesh approach is used by the OpenGGCM (Raeder et al. 2008) and LFM (Lyon et al. 2004) global simulations of the Earth's magnetosphere. The 8-wave solver is one of several methods available in the Space Weather Modeling Framework (SWMF), which has been used for a variety of problems throughout the heliosphere (Toth et al. 2005).

Huba (2005) presents an excellent discussion of the effects of including the Hall term in the MHD equations and the numerical techniques needed to solve them. In the notation of this chapter the inclusion of the Hall term in the generalized Ohm's law results in changes to the energy and induction equations,

$$\frac{\partial \mathcal{E}}{\partial t} + \nabla \cdot \left(\vec{u} \left(\mathcal{E} + P + \frac{B^2}{2\mu_o} \right) - \vec{u} \cdot \frac{\vec{B}\vec{B}}{\mu_o} \right) + \nabla \cdot \left(\vec{u}_H \frac{B^2}{2\mu_o} - 2\frac{1}{\mu_o} \vec{B} (\vec{u}_H \cdot \vec{B}) \right) = 0,$$
(58)

$$\frac{\partial \vec{B}}{\partial t} + \nabla \cdot \left((\vec{u} + \vec{u}_H) \vec{B} - \vec{B} (\vec{u} + \vec{u}_H) \right) = 0, \tag{59}$$

where the "Hall velocity",

$$\vec{u}_H = -\frac{\vec{J}}{ne},\tag{60}$$

has been introduced to clearly illustrate how the Hall terms enter the system of equations. Since these terms are only present in the energy and induction equations it should be clear that the Hall term only transports the magnetic field and energy. To be clear, this means that the Hall effects are not a transport mechanism for mass or momentum. The inclusion of the Hall term introduces a new wave mode, the whistler mode, into the dynamics of the system. The whistler wave speed is significantly larger than the Alfvén speed. This introduces challenges into numerical computation. Since it is the largest wave speed that governs the time step that can be taken within a numerical solution this limitation can result in significant increases in the computational time to the solution. This can be addressed by sub-cycling the Hall physics on the shorter timescale and calculating the ideal MHD physics on the longer timescale.

Of course, the plasma in the Earth's magnetotail and other plasmas throughout the heliosphere can contain more than one ion species so it is often necessary to utilize the multi fluid formulations of the MHD equations to simulate these plasmas. In the notation of this paper these equations are:

$$\frac{\partial \rho_{\alpha}}{\partial t} + \nabla \cdot \rho_{\alpha} \vec{u}_{\alpha} = 0, \tag{61}$$

$$\frac{\partial \rho_{\alpha} \vec{u}_{\alpha}}{\partial t} + \nabla (\rho_{\alpha} \vec{u}_{\alpha} \vec{u}_{\alpha} + I P_{\alpha}) = n_{\alpha} q_{\alpha} (\vec{u}_{\alpha} - \vec{u}_{M}) \times \vec{B} + \frac{n_{\alpha} q_{\alpha}}{n_{e} e} (\vec{J} \times \vec{B} - \nabla P_{e}),$$
(62)

$$\frac{\partial \mathcal{E}_{\alpha}}{\partial t} + \nabla \cdot \left[(\mathcal{E}_{\alpha} + P_{\alpha}) \vec{u}_{\alpha} \right] = \left[n_{\alpha} q_{\alpha} (\vec{u}_{\alpha} - \vec{u}_{M}) \times \vec{B} + \frac{\rho_{\alpha} q_{\alpha}}{n_{e} e} (\vec{J} \times \vec{B} - \nabla P_{e}) \right], \quad (63)$$

$$\frac{\partial \vec{B}}{\partial t} = \nabla \times (\vec{u}_M \times \vec{B}) \tag{64}$$

where the α subscript has been used for the ion species and the term q_{α} allows for the inclusion of higher charge state ions. Furthermore,

$$\vec{u}_M = \frac{1}{en_e} \sum_{\beta} n_{\beta} q_{\beta} \vec{u}_{\beta}$$
(65)

is the charge averaged ion velocity and

$$\vec{J} = en_e(\vec{u}_M - \vec{u}_e) \tag{66}$$

is the current density.

For the electrons, the quasi-neutrality assumption gives,

$$ne = \sum_{\beta} n\rho, \tag{67}$$

as the electron density. Using the definition of current density presented in Eq. (66) we can obtain the electron velocity. The standard fluid equation,

$$\frac{\partial P_e}{\partial t} = -\gamma \nabla (P_e \vec{u}_e) + (\gamma - 1) \vec{u}_e \nabla \cdot P_e, \tag{68}$$

is used to solve for the electron pressure. As this formulation illustrates it is not mathematically possible to cast the multifluid equations in a purely conservative formulation. Numerical techniques used for single fluid have to be adjusted to deal with this situation (Toth et al. 2012 discuss these issues in more detail). It is also worth noting that the energy equation is only true for the hydrodynamic energy density and not the total energy density. In this system to lowest order all the species move in the perpendicular directions with the $E \times B$ velocity. As the magnetic field changes momentum can be transferred between the species in the plasma.

4.2 Incorporation of Internal Plasma Sources in Global MHD Models

In addition to the solar wind plasma, there are various other sources of plasma present in planetary magnetospheres. Plasma sources internal to a planetary magnetosphere may come from the atmosphere/ionosphere, such as the ionospheric outflows at Earth (Chappell 2015;

Welling et al. 2015, this issue) and the planetary ions produced from the exosphere at Mercury (Raines et al. 2015, this issue). In addition, plasma sources may originate from planetary moons and this is especially the case for the gas giants, Jupiter (Bolton et al. 2015, this issue) and Saturn (Blanc et al. 2015, this issue). Through processes like surface warming, active plumes or surface sputtering by magnetospheric particles, moons of the giant planets may possess significant sources of neutrals. The neutrals originating from the moons can become charged particles through various mass-loading processes, thereby supplying plasma to their parent magnetospheres. It is now well known that Io and Enceladus are the major plasma sources of the magnetospheres of Jupiter and Saturn, respectively. The presence of the internal plasma sources to some degree modifies the plasma distribution and composition within the magnetosphere, and in some cases can significantly affect the configuration and dynamics of the magnetosphere. It is, therefore, important to include the internal plasma sources in modeling the structure and dynamics of planetary magnetospheres. Here we provide an overview of the various approaches adopted to incorporate internal plasma sources in global MHD models.

4.2.1 Impact of Ionospheric Outflows

The Alfvén speed in the high-latitude, low-altitude region above the ionosphere is usually very high. Therefore, including this part of the magnetosphere in global magnetosphere simulations imposes severe constraints on the allowable time step that can be used in numerically solving the MHD equations. As a result, presently most global magnetosphere models exclude this region ("gap region") by placing their simulation inner boundaries at altitudes between a couple of and several planetary radii. The ionosphere is conventionally modeled in a separate module as a two-dimensional spherical surface where the electric potential (thus the electric field) is solved for a given distribution of height-integrated conductivity and field-aligned currents (FACs). The FACs are obtained directly from the MHD model of the magnetosphere by first calculating the currents at or near the simulation inner boundary and then mapping them along the dipole field line down to the ionosphere. The electric field obtained from the ionosphere solver is mapped back along the field lines to the magnetosphere boundary, where the $E \times B$ drift velocity is calculated and used to set the boundary condition for plasma velocity. Given the way in which the coupling between the magnetosphere and the ionosphere is treated in present global magnetosphere models, physical processes responsible for producing the ionospheric outflows usually are not directly included in those models. In such cases, the introduction of ionospheric plasma into magnetosphere simulations typically is enabled through prescription of boundary conditions at the low-altitude boundary of the magnetosphere model, similar to the way in which the solar wind plasma is injected into the simulation domain at the sunward boundary. It is worth noting that this type of treatment does not require significant modifications to the MHD equations and is, therefore, relatively convenient in terms of numerical implementation.

Several different approaches have been adopted for adding ionospheric outflows in global MHD models. A relatively simple method is to set the plasma density to relatively high values at the inner boundary and fix it throughout a simulation run. For example, the multifluid MHD model by Winglee et al. (2002) specified constant densities for the light (H^+) and heavy ionospheric species (O^+) at their simulation inner boundary. Pressure gradients and/or other effects (e.g., centrifugal acceleration and numerical diffusion) may drive the ionospheric plasma to flow from the low-altitude boundary into the magnetosphere domain. As such, the ionospheric plasma is added in the simulation in a passive manner in that the outflow parameters are not explicitly set and controlled.

In contrast to the passive method described above, some global models used methods in which the outflow parameters, such as the source location, outflow density and velocity, are explicitly specified at the low-altitude boundary of the magnetosphere model. Several global modeling studies (e.g., Wiltberger et al. 2010; Garcia et al. 2010; Yu and Ridley 2013) performed controlled global simulations to examine the effects of the outflow source location and intensity on the global magnetospheric configuration and dynamics. In these studies, ion outflows were introduced in localized regions, such as the dayside cusp or the nightside auroral zone, and the outflow rates were specified by setting the plasma density and parallel velocity in the boundary conditions.

The choice of outflow parameters may also be made based on empirical outflow models. For example, Brambles et al. (2010) incorporated in the LFM global simulation a driven outflow model based on the empirical model by Strangeway et al. (2005), which was built upon the FAST satellite observations. The empirical model provides a scaling relation between the average outflow flux and the average earthward-flowing Poynting flux, which is calculated directly from the MHD model near the inner boundary. This approach in effect enables a two-way coupling between the magnetosphere and the ionosphere, because the outflow source location and intensity may vary in time depending on the magnetospheric conditions.

More self-consistent implementation of ionospheric outflows may be achieved by coupling a global MHD model with a physics-based ionospheric outflow model. Glocer et al. (2009) coupled the Polar Wind Outflow Model (PWOM) into the SWMF to study the effects of polar wind type outflows on the coupled magnetosphere-ionosphere system. PWOM includes important physical processes responsible for the transport and acceleration of the ionospheric gap region between the magnetosphere and ionosphere. It takes inputs from both the magnetosphere model (FACs and plasma convection pattern) and the upper atmosphere model (neutral densities and neutral winds) to calculate the upwelling and outflowing of ionospheric plasma. In return, the outflow fluxes obtained at the top boundary of the PWOM model are used to set the inner boundary conditions of the magnetosphere model.

4.2.2 Plasma Sources Associated with Planetary Satellites

Different from the Earth's magnetosphere where the magnetospheric plasma comes either from the solar wind or the ionosphere, the bulk of the magnetospheric plasma in the giant planet magnetospheres originate predominantly from planetary satellites. At Jupiter, the major plasma source is the volcanic moon, Io, which supplies plasmas to the Jovian magnetosphere at a rate of 260–1400 kg/s (Bagenal and Delamere 2011). At Saturn, the dominant source of magnetospheric plasma is the icy moon, Enceladus, which produces predominantly water-group ions to the magnetosphere at a rate of 12–250 kg/s (Bagenal and Delamere 2011). At both planets, the presence of internal plasma sources plays a crucial role in shaping the magnetosphere. It is, therefore, essential to include the internal plasma sources associated with the moons in global models of the giant planet magnetospheres.

There are, in general, two types of approaches used for incorporating plasma sources associated with moons. One relies on prescription of boundary conditions, similar to the approach outlined above for incorporating ionospheric outflows into Earth's magnetosphere models. For example, the global MHD model by Ogino et al. (1998) which was first applied to Jupiter and later adapted to Saturn (Fukazawa et al. 2007a, 2007b), does not explicitly include in the simulation domain plasma sources associated with moons. Rather, the model included the internal plasma sources by fixing plasma density and pressure in

time at the inner boundary, which was placed outside of the main regions in which moonassociated plasmas are added to the systems. Similarly, in the multi-fluid MHD model applied to Saturn's magnetosphere, Kidder et al. (2009) held the densities of various plasma fluids fixed near their simulation inner boundary to mimic the addition of new plasma from Enceladus.

The other approach used in the modeling of the giant planets' magnetospheres incorporates internal plasma sources associated with moons in an explicit manner. The neutral gases emanating from the moons in the Jovian and Saturnian magnetospheres are distributed in a broad region forming plasma and neutral tori, which mass-load newly created charged particles which then modify the plasma flow in the system via electromagnetic forces (see a review by Szegö et al. 2000). This occurs not only near the vicinities of the moons, but also over extended regions of space. It is desirable to self-consistently take into account this effect in a global magnetosphere model. This can be done by incorporating appropriate source and loss terms into the MHD equations described above. One can derive the mass-loading source terms for MHD using first-principles from the Boltzman equation (Cravens 1997; Gombosi 1998). Terms describing the change of the plasma phase-space distribution due to collisional processes, including ionization, charge-exchange, recombination, and elastic collisions, can be included in the Boltzman equation. Appropriate velocity moments can then be taken to obtain the source terms associated with various mass-loading processes for the continuity, momentum and energy equations of MHD. One advantage of this method over the boundary condition method is that it describes in a self-consistent way the change of mass, momentum and energy of magnetospheric plasma due to mass-loading. This approach has been used in global models of the giant planets' magnetospheres, such as the SWMF applications to Saturn's magnetosphere by Hansen et al. (2005), Jia et al. (2012a, 2012b), and Jia and Kivelson (2012) and the global MHD model of Jupiter's magnetosphere by Chané et al. (2013).

4.3 Hybrid Models

The most common hybrid approach used in simulating space plasmas treats the ions kinetically and the electrons as a massless charge neutralizing fluid. In the hybrid regime, the density, temperatures and magnetic field is such that the ions are essentially collisionless. On the other hand the electrons have relatively small gyroradii and may undergo an order of magnitude or more collisions. Thus the electrons are described as a massless collisiondominated thermal fluid. There are finite electron mass hybrid schemes in existence, which will not be discussed here. Hybrid schemes have been around for many years thus the interested reader should see the reviews by Brecht and Thomas (1988), Lipatov (2002), Winske et al. (2003), and the references therein for historical perspectives. The most recent review is that of Ledvina et al. (2008), where the following brief description is taken from.

The hybrid approach starts with the following assumptions.

(i) Quasi-neutrality is assumed,

$$n_e = \sum_i n_i \tag{69}$$

Thus the displacement current is ignored in Ampere's law (Eq. (74)). This assumption is valid on scales larger than the Debye length. The assumption breaks down when the grid resolution is finer than the Debye length. This also implies that $\nabla \cdot J = 0$, and removes most electrostatic instabilities.

(ii) The Darwin approximation is assumed.

This approximation splits the electric field into a longitudinal part E_L and a solenoidal part E_T . Then $\nabla \times E_L = 0$ and $\nabla \cdot E_T = 0$ and $\partial E_T / \partial t$ is neglected in Ampere's law (Eq. (74)). This allows the light waves to be ignored. It also removes relativistic phenomena.

- (iii) The mass of the electrons is taken to be zero.
- (iv) The electrons collectively act as a fluid.

Thus the electron plasma and gyrofrequencies are removed from the calculations. This means that high frequency modes are not present, such as the electron whistler. By using these last two assumptions there is no longer a physical mechanism to describe the system behavior at small scales. The Debye length and the magnetic skin depth are not meaningful in this scheme. This sets the limit on the cell size that should be used to at least an order of magnitude larger than the electron skin depth c/ω_{pe} . It is possible to use cell sizes less than the ion skin depth but the results are meaningless. The chosen cell size is much larger than the kinetic scales all that is accomplished is the creation of the world's most expensive MHD simulation.

With these assumptions the hybrid scheme solves the following ion momentum and position equations for each particle:

$$\frac{dv}{dt} = \frac{q}{m_i} [\boldsymbol{E} + \boldsymbol{v} \times \boldsymbol{B}] - \eta \boldsymbol{J}_{total}$$
(70)

$$\frac{d\boldsymbol{x}}{dt} = \boldsymbol{v} \tag{71}$$

where J is the total current density and η is the plasma resistivity. The electron momentum equation can be written as:

$$\boldsymbol{E} = \frac{1}{n_i e} \Big[(\nabla \times \boldsymbol{B}) \times \boldsymbol{B} - \boldsymbol{J}_i \times \boldsymbol{B} - \nabla (n_e T_e) + \eta \boldsymbol{J}_{total} \Big]$$
(72)

With the electron temperature given by:

$$\frac{\partial T_e}{\partial t} + \boldsymbol{u}_e \cdot \nabla T_e + \frac{3}{2} T_e \cdot \boldsymbol{u}_e = \frac{2}{3n_e} \eta \boldsymbol{J}_{total}^2$$
(73)

Here T_e is the electron temperature and u_e is the electron velocity. Note that (73) does not include the effects of thermal conduction, but that can be added if appropriate. Ampere's law becomes:

$$\nabla \times \boldsymbol{H} = \boldsymbol{J}_i + \boldsymbol{J}_e \tag{74}$$

where J_i and J_e are the ion and electron current densities. The magnetic field is obtained from Faraday's law, given below:

$$\nabla \times \boldsymbol{E} + \frac{\partial \boldsymbol{B}}{\partial t} = 0 \tag{75}$$

The electric field contains contributions from the electron pressure gradient, resistive effects and Hall currents. The scheme correctly simulates electromagnetic plasma modes up to and including the lower portion of the whistler wave spectrum (well below the electron cyclotron frequency, $\omega \ll \omega_{ce}$). Shock formation physics is included, therefore no assumptions

or shock capturing techniques are needed to capture a shock. The time step is determined by the ion cyclotron frequency. This comes at the price of the loss of electron particle effects and charge separation. Some small-scale electrostatic effects can be included through the resistivity terms. The resistivity terms can also be used to stabilize the numerical scheme used to solve the equations by adding it in as a small amount of artificial resistivity.

4.4 Magnetosphere-Ionosphere Coupling

The ionosphere-magnetosphere coupling is not a process in itself. It is rather a chain of processes that act as a control loop between the dynamics of the ionospheric and of the magnetospheric plasmas connected by conductive magnetic field lines as shown in Fig. 8. A modification of the transport in one region has consequences on the transport in the conjugate region and that affects in turn the initial transport in the first region. For example, the convection in the magnetosphere results in convection in the ionosphere (see Fig. 9). The plasma dynamics in one region is constrained by the dynamics in the other. For each region, the ionosphere-magnetosphere coupling could be assimilated to some kind of interactive boundary conditions (representing the interaction with the conjugate region) that need to be solved self-consistently with the dynamics of the region considered.

In a first approach, the ionospheric plasma exhibits local-time, latitudinal, seasonal variations but forms a continuous conductive shell embedded in the high-altitude planetary atmosphere. It lies at the footprints of conductive planetary magnetic field lines that connect it to different magnetospheric regions. The polar cap magnetic field lines are open with one footprint in the polar ionosphere and the other end extended to large distances downtail, in the so-called lobes. The lobe plasma is believed to be diluted and therefore does not develop significant couplings with the ionosphere. Near the equator, the magnetic field lines remain fully embedded in the topside ionosphere and do not reach the magnetosphere. Between the polar cap and the equatorial strip, the magnetic field lines are closed with both footprints in the ionosphere and their apex reach the magnetosphere. Near the planet, a region called "plasmasphere" filled with cold plasma of ionospheric origin in corotation with the planet may exist, as well as radiation belts with very energetic particles trapped on closed orbits around the planet. The so-called "plasmasheet" represents the main plasma reservoir in the magnetospheres of Earth, Jupiter and Saturn. The transport mechanisms differ for each planet: they involve corotation, outward diffusion from inner plasma sources or earthward convection of plasma ultimately extracted from external sources (solar wind), but all result in the formation of a dense and hot plasma sheet, confined near the equatorial plane and extending up to large distances down tail. The conductive magnetic field lines allow electric field transmission, current circulation and particle exchanges. The effects of these magnetic-field-aligned processes are enhanced when they involve dense and dynamical regions such as the ionosphere and the plasmasheet, resulting in significant consequences on the dynamics of both regions at large scales as well as at local or transient scales.

The coupled ionosphere-magnetosphere system can be described by a feedback loop derived from various investigations in the terrestrial environment (Vasyliunas 1970; Wolf 1979; Harel et al. 1981; Fontaine et al. 1985; Peymirat and Fontaine 1994) as illustrated in Fig. 13, where the magnetospheric plasma is indicated in the top row. External sources such as the planetary rotation or the solar wind-magnetosphere dynamo contribute to produce large-scale electric fields in the magnetospheric plasma into a large-scale motion. Smaller-scale processes, instabilities, phase space diffusion processes, etc. add smaller-scale motions and contribute to the global and local plasma distribution and current circulation in the magnetosphere.



Fig. 16 Feedback loop illustrating the ionosphere-magnetosphere coupling with magnetospheric (*red*), ionospheric (*blue*) and field-aligned (*green*) electrodynamic parameters and processes. *Black rectangles* represent external sources

Field-aligned processes are shown in the second row of Fig. 16. On one hand, the current closure $\nabla \cdot \mathbf{j}_M = 0$ in the magnetosphere, where \mathbf{j}_M is the magnetospheric current density, implies a current circulation along magnetic field lines \mathbf{j}_{\parallel} down to the ionosphere. On the other hand, particles with pitch-angles smaller than the atmospheric loss cone reach the ionosphere at the footprint of magnetic field lines: they contribute to the field-aligned currents. The mirror effect due to the magnetic field line convergence limits the particle fluxes that reach the ionosphere and thus the field-aligned current density transmitted to the ionosphere. Current-voltage relations, such as those proposed by Knight (1973) (see Eqs. (39) and (40)), predict that parallel potentials can develop and increase the field-aligned current density when the available precipitating fluxes cannot match the current density required by the current closure in the magnetosphere. Ionospheric particles can also escape from the ionosphere, in particular electrons which are very mobile along magnetic field lines. They carry return currents due to a favorable effect of the mirror force from the ionosphere toward the magnetosphere. It is generally difficult to measure particle outflows of ionospheric origin due to their low energy, except if they are accelerated (see Chappell 2015).

The UV and EUV solar radiation contribute to create an ionospheric layer in the highaltitude atmosphere. The dynamics of the ionosphere is governed by the ionospheric Ohm's law:

$$\boldsymbol{j}_{I} = \boldsymbol{\sigma}(\boldsymbol{E} + \boldsymbol{V}_{n} \times \boldsymbol{B}) \tag{76}$$

and the ionospheric current closure equation:

$$\nabla \cdot \boldsymbol{j}_I = 0 \tag{77}$$

where j_I is the ionospheric current density, σ the ionospheric conductivity tensor, E_I the electric field at ionospheric altitudes, V_n the velocity of the neutral wind, B the magnetic field. In addition of this solar source, the fluxes of energetic magnetospheric precipitating particles into the ionosphere contribute to produce the well-known auroral light emissions and also ionization. The resulting conductivity enhancements and the presence of field-aligned currents modify the distribution of perpendicular electric currents and electric fields at the ionosphere via magnetic field lines by taking into account the eventual presence of parallel electric fields. This new electric field distribution modifies in turn the plasma transport in the magnetosphere, which closes the feedback loop.

Finally, any modification/event at large or smaller scales that occurs in one region is transmitted to the other one where it modifies its own dynamics. However, the possibilities of exchanges of particles, momentum, and energy are limited by the plasma configuration in each region. A mismatch between both regions can be overcome by the set up of field-aligned electric fields and currents, in the limit of energy density available in each region. These effects result in parallel particle acceleration and thus in light emissions when accelerated particles precipitate into the ionosphere/upper atmosphere.

4.4.1 Time-Varying Coupling

The above description does not only apply to quasi-steady ionosphere-magnetosphere coupling, but works similarly at smaller-scales (see Lysak 1990 for a review). For example, time-varying fluctuations in the magnetosphere or wave-particle interactions occurring during plasma transport may generate Alfvén waves that carry field-aligned currents. These currents close similarly through the ionosphere. They result in fluctuating effects in the ionosphere that will affect auroras, conductivities, electric fields and currents. Fluctuating conditions in the ionosphere are in turn transmitted to the magnetosphere via magnetic field lines and produce fluctuating feedback effects. The superposition of initial and feedback fluctuations can stabilize or destabilize the plasma; it can also give rise to periodic effects as pulsations, formation of multiple arcs, etc.

Small-scale processes such as magnetic reconnection imply a connectivity interruption and reconfiguration for a subset of magnetic field lines in a localized region. On the reconnection time scale, field-aligned processes cannot exist because of connectivity changes and the ionosphere and magnetosphere dynamics are disconnected. This is not the case for the time periods just before and after reconnection: important effects occur in both regions, resulting in enhanced field-aligned couplings, i.e. large field-aligned particle fluxes, electric fields and currents.

4.4.2 Planet-Moon Interactions

The interaction of magnetized planets with moons is another example of local feedback processes. It depends on the electrical properties of the moons, or rather of the obstacle, and on the flow characteristics (for a review, see for example Kivelson et al. 2004). The obstacle can be the magnetic field, the atmosphere and ionosphere or the body itself depending of the radial variation of the energy density. The magnetospheric flow is coupled to the planetary ionosphere via magnetic field lines and this coupling drags the magnetospheric plasma at a speed which may differ from the moons' orbital velocity. If the flow velocity in the rest frame of the moon were super-Alfvénic, it would produce a shock wave ahead of the obstacle as in

the solar wind/magnetosphere interaction. Inside magnetospheres, the interaction velocity is usually sub-Alfvénic.

In the case of an insulating body, the sub-Alfvénic magnetospheric flow is absorbed by the surface, an initially empty wake appears downstream and the magnetic field exhibits only weak perturbations. Ions can be created from various interaction processes between the magnetospheric particles and the moon, and this so-called ion pickup source contributes to the mass-loading of the magnetospheric flow.

In the case of a conducting body, the sub-Alfvénic magnetospheric flow slows upstream of the body, the planetary magnetic field lines get bent and shear Alfvén waves are launched. These waves carry field-aligned currents and they generate perturbations in the field which are known as Alfvén wings. Alfvén wings form an angle Θ_A with the initial magnetospheric magnetic field:

$$\theta_A = \tan^{-1} \left(\frac{V_M}{V_A} \right) \tag{78}$$

where V_M is the velocity of the magnetospheric flow, and V_A the Alfvén velocity.

They extend down to the planetary ionosphere which allows the current closure. This localized ionosphere-magnetosphere coupling contributes to divert the magnetospheric plasma flow around the conducting body and all along the Alfvén wings. It modifies locally the properties in the magnetically conjugated ionosphere. For example, light emissions in the ionosphere at the magnetic footprints of the Galilean moons in the Jovian magnetosphere represent the signature of this localized ionosphere-magnetosphere coupling.

In the case of a magnetized body, the moon's magnetic field creates a small magnetosphere inside the planetary magnetosphere. Up to now, Ganymede is the only known magnetized moon in the solar system. Although very small, Ganymede's magnetosphere contains features similar to terrestrial and planetary magnetospheres (e.g., Kivelson et al. 1998), as the presence of a magnetopause, innermost regions protected by the internal magnetic field, and auroras (Jia et al. 2009). One of the differences is that polar magnetic field lines from Ganymede's polar region connect the Jovian ionosphere at their other end. They carry fieldaligned currents and contribute to a local coupling between the planetary ionosphere and the moon's magnetosphere embedded in the magnetospheric flow.

5 Summary

In this paper, the basic and common processes, related to plasma supply to each region of the planetary magnetospheres in our solar system, were reviewed. In addition to major processes related to the source, transport, energization, loss of the magnetospheric plasmas, basic equations and modeling methods, with a focus on plasma supply processes for planetary magnetospheres, are also reviewed. The topics reviewed in this paper can be summarized as follows: Source Processes related to the surface (Sect. 1.1), ionosphere (Sect. 1.2), and solar wind (Sect. 1.3). Section 2 is dedicated to processes related to the transport and energization of plasma such as Axford/Hines cycle (Sect. 2.1), Dungey cycle (Sect. 2.2), rotational driven transport and Vasyliunas cycle (Sect. 2.3), field-aligned potential drop (Sect. 2.4), non-adiabatic acceleration (Sect. 2.5), and pick-up acceleration and mass loading (Sect. 2.6). In Sect. 3. loss processes related to the tail reconnection and plasmoids (Sect. 3.1), charge exchange (Sect. 3.2), and precipitations into planets (Sect. 3.3) are reviewed. Section 4 contains an overview of basic equations and modeling methods, which includes MHD simulation

(Sect. 4.1), incorporation of internal plasma sources in global MHD models (Sect. 4.2), hybrid models (Sect. 4.3), and magnetosphere-ionosphere coupling (Sect. 4.4). The review provides the basic knowledge to understand various phenomena in planetary magnetospheres described in the following chapters.

Acknowledgements This work was partially supported by Grant-In-Aid for Scientific Research (B) 24340118 from JSPS and MEXT of Japan. KS also thanks for supports from the GEMSIS project at STEL, Nagoya University. CMJ's work at Southampton was supported by a Royal Astronomical Society Fellowship and a Science and Technology Facilities Council Ernest Rutherford Fellowship. FC is funded by JPL contract 1467206. MW is supported by NASA grants NNH12AU10I and NNH11AR82I. The National Center for Atmospheric Research is sponsored by the National Science Foundation. XJ is supported by NASA grants NNX12AK34G and NNX12AM74G, and by NSF grant AGS 1203232.

Ethical Statement This manuscript is prepared to submit to SSR as a review article after discussion at the ISSI workshop in 2013 and never submitted to elsewhere. Contents of this manuscript have nothing to do with the following issues:

- Disclosure of potential conflicts of interest,
- · Research involving Human Participants and/or Animals,
- Informed consent.

References

- T. Abe et al., Observations of polar wind and thermal ion outflow by Akebono/SMS. J. Geomagn. Geoelectr. **48**, 319 (1996)
- H. Andersen, H.L. Bay, in *Sputtering by Particle Bombardment I*, ed. by R. Behrisch (Springer, Berlin, 1981), Chap. 4
- M. Andre, A.L. Yau, Theories and observations of ion energization and outflow in the high latitude magnetosphere. Space Sci. Rev. 80, 27 (1997)
- V.G. Anicich, Evaluated bimolecular ion-molecule gas phase kinetics of positive ions for use in modeling planetary atmospheres, cometary comae, and interstellar clouds. J. Phys. Chem. Ref. Data 22, 1469– 1569 (1993)
- M. Ashour-Abdalla et al., The formation of the wall region—Consequences in the near-Earth magnetotail. Geophys. Res. Lett. 19, 1739 (1992)
- M. Ashour-Abdalla et al., Consequences of magnetotail ion dynamics. J. Geophys. Res. 99, 14891 (1994)
- W.I. Axford, Viscous interaction between the solar wind and the Earth's magnetosphere. Planet. Space Sci. 12, 45 (1964)
- W.I. Axford, C.O. Hines, A unifying theory of high-latitude geophysical phenomena and geomagnetic storms. Can. J. Phys. 39, 1433 (1961)
- S.V. Badman, S.W.H. Cowley, Significance of Dungey-cycle flows in Jupiter's and Saturn's magnetospheres, and their identification on closed equatorial field lines. Ann. Geophys. 25, 94 (2007)
- F. Bagenal, The magnetosphere of Jupiter: Coupling the equator to the poles. J. Atmos. Sol.-Terr. Phys. 69, 387 (2007). doi:10.1016/j.jastp.2006.08.012
- F. Bagenal, P.A. Delamere, Flow of mass and energy in the magnetospheres of Jupiter and Saturn. J. Geophys. Res. 116, A05209 (2011). doi:10.1029/2010JA016294
- F. Bagenal, J.D. Sullivan, Direct plasma measurements in the Io torus and inner magnetosphere of Jupiter. J. Geophys. Res. 86, 8447 (1981)
- J. Bailey, M. Gruntman, Experimental study of exospheric hydrogen atom distributions by Lymanalpha detectors on the TWINS mission. J. Geophys. Res. (Space Phys.) 116(A15), 302 (2011). doi:10.1029/2011JA016531
- P.M. Banks, T.E. Holzer, High-latitude plasma transport: The polar wind. J. Geophys. Res. 74, 6317 (1969). doi:10.1029/JA074i026p06317
- R.A. Baragiola et al., Nucl. Instrum. Methods Phys. Res. B 209, 294 (2003)
- L.F. Bargatze, D.N. Baker, R.L. McPherron, E.W. Hones, Magnetospheric impulse response for many levels of geomagnetic activity. J. Geophys. Res. 90, 6387 (1985). doi:10.1029/JA090iA07p06387
- R. Behrisch, W. Eckstein, Sputtering by Particle Bombardment: Experiments and Computer Calculations from Threshold to MeV Energies (Springer, Berlin, 2007)
- A. Bhardwaj, G.R. Gladstone, Auroral emissions of the giant planets. Rev. Geophys. 38, 295 (2000)

- J.P. Biersack, W. Eckstein, Sputtering of solids with the Monte Carlo program TRIM.SP. Appl. Phys. A 34, 73 (1984)
- G.A. Bird, Molecular Gas Dynamics and the Direct Simulation of Gas Flows (Clarendon Press, Oxford, 1994)
- K. Birkeland, Sur la déviabilité magnétique des rayons corpusculaires provenant du Soleil. C. R. Acad. Sci. 150, 246 (1910)
- J. Bishop, Multiple charge exchange and ionization collisions within the ring current-geocoronaplasmasphere system: generation of a secondary ring current on inner L shells. J. Geophys. Res. 101, 17,325 (1996). doi:10.1029/95JA03468
- M. Blanc et al., Space Sci. Rev. (2015, this issue)
- S.J. Bolton et al., Space Sci. Rev. (2015, this issue)
- S.J. Bolton, Jupiter's inner radiation belts, in Jupiter: The Planet, Satellites, and Magnetosphere, ed. by F. Bagenal et al. (Cambridge Univ. Press, Cambridge, 2004), p. 671, Chap. 27
- S.J. Bolton, R.M. Thorne, S. Bourdarie, I. de Pater, B. Mauk, Jupiter's inner radiation belts, in *Jupiter: The Planet, Satellites, and Magnetosphere*, ed. by F. Bagenal et al.(Cambridge Univ. Press, Cambridge, 2004), p. 671. Chapter 27
- B. Bonfond, When Moons create Aurora: the satellite footprints on giant planets, in Auroral Phenomenology and Magnetospheric Processes: Earth and Other Planets. AGU Geophysical Monograph Series, vol. 197 (2012), p. 133
- J.W. Boring et al., Sputtering of solid SO₂. Nucl. Instrum. Methods B 1, 321 (1984)
- J.E. Borovsky, M.F. Thomnsen, R.C. Elphic, The driving of the plasma sheet by the solar wind. J. Geophys. Res. 103(A8), 17,617 (1998). doi:10.1029/97JA02986
- J.E. Borovsky et al., What determines the reconnection rate at the dayside magnetosphere? J. Geophys. Res. 113, A07210 (2008). doi:10.1029/2007JA012645
- O.J. Brambles et al., Effects of causally driven cusp O⁺ outflow on the storm time magnetosphereionosphere system using a multifluid global simulation. J. Geophys. Res. **115**, A00J04 (2010). doi:10.1029/2010JA015469
- S.H. Brecht, V.A. Thomas, Multidimensional simulations using hybrid particle codes. Comput. Phys. Commun. 48, 135–143 (1988)
- N.M. Brice, G.A. Ioannidis, The magnetospheres of Jupiter and Earth. Icarus 13, 173 (1970)
- W.L. Brown, R.E. Johnson, Sputtering of ices: a review. Nucl. Instrum. Methods B 13, 295 (1986)
- W.L. Brown, W.M. Augustyniak, K.J. Marcantonio, E.N. Simmons, J.W. Boring, R.E. Johnson, C.T. Reimann, Electronic sputtering of low temperature molecular solids. Nucl. Instrum. Methods Phys. Res. B 1, 307 (1984)
- D.G. Brown, J.L. Horwitz, G.R. Wilson, Synergistic effects of hot plasma-driven potentials and wave-driven ion heating on auroral ionospheric plasma transport. J. Geophys. Res. 100, 17,499 (1995)
- M. Bruno, G. Cremonese, S. Marchi, Neutral sodium atoms release from the surfaces of the Moon and Mercury induced by meteoroid impacts. Planet. Space Sci. 55, 1494 (2007)
- J. Büchner, L.M. Zelenyi, Regular and chaotic charged particle motion in magnetotaillike field reversals: 1. Basic theory of trapped motion. J. Geophys. Res. 94, 11,821 (1989)
- J.L. Burch et al., Properties of local plasma injections in Saturn's magnetosphere. Geophys. Res. Lett. 32, L14S02 (2005). doi:10.1029/2005GRL022611
- L.F. Burlaga, Magnetic fields and plasmas in the inner heliosphere: Helios results. Planet. Space Sci. 49, 1619 (2001)
- R.K. Burton, R.L. McPherron, C.T. Russell, Terrestrial magnetosphere—half-wave rectifier of interplanetary electric-field. Science 189(4204), 717 (1975a)
- R.K. Burton, R.L. McPherron, C.T. Russell, An empirical relationship between interplanetary conditions and Dst. J. Geophys. Res. 80, 4204 (1975b)
- M.N. Caan, R.L. McPherron, C.T. Russell, Solar wind and substorm-related changes in the lobes of the geomagnetic tail. J. Geophys. Res. 78, 8087 (1973)
- C.W. Carlson, R.F. Pfaff, J.G. Watzin, Fast Auroral Snapshot (FAST) mission. Geophys. Res. Lett. 25, 2013 (1998)
- T.A. Cassidy, R.E. Johnson, Monte Carlo model of sputtering and other ejection processes within a regolith. Icarus 176, 499 (2005)
- T.A. Cassidy et al., The spatial morphology of Europa's near-surface O₂ atmosphere. Icarus **191**, 755 (2007)
- T.A. Cassidy et al., Radiolysis and photolysis of icy satellite surfaces: experiments and theory. Space Sci. Rev. 153(1–4), 299 (2010)
- T.A. Cassidy et al., Magnetospheric ion sputtering and water ice grain size at Europa. Planet. Space Sci. 77, 64 (2013)
- B.L. Cecconi et al., Goniopolarimetric study of the Rev 29 perikrone using the Cassini/RPWS/HFR radio receiver. J. Geophys. Res. 114, A03215 (2009)

- J.W. Chamberlain, Planetary coronae and atmospheric evaporation. Planet. Space Sci. 11, 901 (1963). doi:10.1016/0032-0633(63)90122-3
- E. Chané, J. Saur, S. Poedts, Modeling Jupiter's magnetosphere: Influence of the internal sources. J. Geophys. Res. 118, 2157 (2013). doi:10.1002/jgra.50258
- C.R. Chappell, The role of the ionosphere in providing plasma to the terrestrial magnetosphere—an historical overview. Space Sci. Rev. (2015, in press)
- F.F. Chen, Introduction to Plasma Physics and Controlled Fusion. Boom Koninklijke Uitgevers (1984)
- J. Chen, P.J. Palmadesso, Chaos and nonlinear dynamics of single-particle orbits in magnetotaillike magnetic field. J. Geophys. Res. 91, 1499 (1986)
- Y.T. Chiu, M. Schulz, Slf-consistent particle and parallel electrostatic electric field distributions in the magnetospheric-ionospheric auroral region. J. Geophys. Res. 83, 629 (1978)
- M.J. Cintala, Impact induced thermal effects in the lunar and Mercurian regoliths. J. Geophys. Res. 97, 947 (1992)
- J.B. Cladis, Parallel acceleration and transport of ions from polar ionosphere to plasma sheet. Geophys. Res. Lett. 13, 893 (1986)
- J.T. Clarke, Auroral processes on Jupiter and Saturn, in Auroral Phenomenology and Magnetospheric Processes: Earth and Other Planets. AGU Geophysical Monograph Series, vol. 197 (2012), p. 113
- J.E.P. Connerney, J.H. Waite, New model of Saturn's ionosphere with an influx of water. Nature **312**, 136 (1984)
- J.E.P. Connerney et al., Images of excited H⁺₃ at the foot of the Io flux tube in Jupiter's atmosphere. Science **262**, 1035–1038 (1993)
- J.F. Cooper et al., Energetic ion and electron irradiation of the icy Galilean satellites. Icarus 149, 133 (2001)
- S.W.H. Cowley, The causes of convection in the Earth's magnetosphere: A review of developments during the IMS. Rev. Geophys. 20(3), 531–565 (1982). doi:10.1029/RG020i003p00531
- S.W.H. Cowley, E.J. Bunce, R. Prangé, Saturn's polar ionospheric flows and their relation to the main auroral oval. Ann. Geophys. 22, 1379 (2004)
- T.E. Cravens, *Physics of Solar System Plasmas* (Camb. Univ. Press, Cambridge, 1997). doi:10.1017/ CBO9780511529467
- J. De Keyser, R. Maggiolo, M. Echim, Monopolar and bipolar auroral electric fields and their effects. Ann. Geophys. 28, 2027 (2010)
- A.E. De Vries et al., Synthesis and sputtering of newly formed molecules by kiloelectronvolt ions. J. Phys. Chem. 88, 4510 (1984)
- P.A. Delamere, F. Bagenal, Solar wind interaction with Jupiter 's magnetosphere. J. Geophys. Res. 115, A10201 (2010). doi:10.1029/2010JA015347
- P.A. Delamere et al., Magnetic signatures of Kelvin-Helmholtz vortices on Saturn's magnetopause: Global survey. J. Geophys. Res. 118, 393 (2013)
- D.C. Delcourt, Particle acceleration by inductive electric fields in the inner magnetosphere. J. Atmos. Sol.-Terr. Phys. 64, 551 (2002)
- D.C. Delcourt, J.-A. Sauvaud, Populating of the cusp and boundary layers by energetic (hundreds of keV) equatorial particles. J. Geophys. Res. 104, 22,635 (1999)
- D.C. Delcourt, J.-A. Sauvaud, A. Pedersen, Dynamics of single-particle orbits during substorm expansion phase. J. Geophys. Res. 95, 20,853 (1990)
- D.C. Delcourt, R.F. Martin Jr., F. Alem, A simple model of magnetic moment scattering in a field reversal. Geophys. Res. Lett. 21, 1543 (1994)
- D.C. Delcourt et al., On the nonadiabatic precipitation of ions from the near-Earth plasma sheet. J. Geophys. Res. 101, 17,409 (1996)
- J. Dessler, E.N. Parker, Hydromagnetic theory of geomagnetic storms. J. Geophys. Res. 64, 2239 (1959). doi:10.1029/JZ064i012p02239
- G.A. DiBraccio, J.A. Slavin, S.A. Boardsen, B.J. Anderson, H. Korth, T.H. Zurbuchen, J.M. Raines, D.N. Baker, R.L. McNutt Jr., S.C. Solomon, MESSENGER observations of magnetopause structure and dynamics at Mercury. J. Geophys. Res. 118, 997 (2013). doi:10.1002/jgra.50123
- J.W. Dungey, Interplanetaly magnetic field and the auroral zones. Phys. Rev. Lett. 6, 47 (1961)
- W. Eckstein, H.M. Urbassek, Computer simulation of the sputtering process, in *Sputtering by Particle Bombardment: Experiments and Computer Calculations from Threshold to MeV Energies*, ed. by R. Behrisch, W. Eckstein (Springer, Berlin, 2007), p. 21
- G. Eichhorn, Heating and vaporization during hypervelocity particle impact. Planet. Space Sci. 26, 463 (1978a)
- G. Eichhorn, Primary Velocity Dependence of impact ejecta parameters. Planet. Space Sci. 26, 469 (1978b)
- R.E. Ergun et al., S bursts and the Jupiter ionospheric Alfvén resonator. J. Geophys. Res. 111, A06212 (2006)

- D.S. Evans, Precipitation electron fluxes formed by magnetic-field-aligned potential differences. J. Geophys. Res. 79, 2853 (1974)
- H.J. Fahr, The extraterrestrial UV-background and the nearby interstellar medium. Space Sci. Rev. 15, 483 (1974). doi:10.1007/BF00178217
- M. Famà, J. Shi, R.A. Baragiola, Sputtering of ice by low-energy ions. Surf. Sci. 602, 156 (2008)
- C.J. Farrugia, N.V. Erkaev, H.K. Biernat et al., Anomalous magnetosheath properties during Earth passage of an interplanetary magnetic cloud. J. Geophys. Res. 100, 19245 (1995)
- Y.I. Feldstein et al., Structure of the auroral precipitation region in the dawn sector: relationship to convection reversal boundaries and field-aligned currents. Ann. Geophys. 19, 495 (2001)
- W.L. Fite, T.R. Brackman, W.R. Snow, Charge transfer in proton-hydrogen atom collisions. Phys. Rev. 112, 1161 (1958)
- M.-C. Fok et al., J. Geophys. Res. 111 (2006). doi:10.1029/2006JA011839
- D. Fontaine et al., Numerical simulation of the magnetospheric convection including the effect of electron precipitation. J. Geophys. Res. 90, 8343 (1985)
- R.J. Forsyth et al., The underlying Parker spiral structure in the Ulysses magnetic field observations, 1990– 1994. J. Geophys. Res. 101, 395 (1996)
- J.L. Fox, K.Y. Sung, Solar activity variations of the Venus thermosphere/ionosphere. J. Geophys. Res. 106, 21305 (2001)
- L.A. Frank, On the extraterrestrial ring current during geomagnetic storm. J. Geophys. Res. 72, 3753 (1967)
- K.M. Frederick-Frost et al., SERSIO: Svalbard EISCAT rocket study of ion outflows. J. Geophys. Res. 112, A08307 (2007). doi:10.1029/2006JA011942
- K. Fukazawa, T. Ogino, R.J. Walker, Magnetospheric convection at Saturn as a function of IMF Bz. Geophys. Res. Lett. 34(1) (2007a). doi:10.1029/2006GL028373
- K. Fukazawa, T. Ogino, R.J. Walker, Vortex-associated reconnection for northward IMF in the Kronian magnetosphere. Geophys. Res. Lett. 34 (2007b). doi:10.1029/2007GL031784
- S.A. Fuselier, W.S. Lewis, Properties of near-earth magnetic reconnection from in-situ observations. Space Sci. Rev. 160(1–4), 95 (2011)
- S.A. Fuselier et al., Energetic neutral atoms from the Earth's subsolar magnetopause. Geophys. Res. Lett. 371, 101 (2010). doi:10.1029/2010GL044140
- P. Galopeau, P. Zarka, D. Le Quéau, Source location of SKR: the Kelvin-Helmholtz instability hypothesis. J. Geophys. Res. 100, 26397 (1995)
- S.B. Ganguli, The polar wind. Rev. Geophys. 34, 311 (1996)
- K.S. Garcia, V.G. Merkin, W.J. Hughes, Effects of nightside O⁺ outflow on magnetospheric dynamics: Results of multifluid MHD modeling. J. Geophys. Res. 115, A00J09 (2010). doi:10.1029/2010JA015730
- D.J. Gershman, J.A. Slavin, J.M. Raines et al., Magnetic flux pileup and plasma depletion in Mercury's subsolar magnetosheath. J. Geophys. Res. 118, 7181 (2013)
- A. Glocer, G. Tóth, T. Gombosi, D. Welling, Modeling ionospheric outflows and their impact on the magnetosphere, initial results. J. Geophys. Res. 114(A), 05,216 (2009). doi:10.1029/2009JA014053
- T.I. Gombosi, Physics of the Space Environment (Cambridge University Press, Cambridge, 1998)
- J.T. Gosling et al., The electron edge of the low latitude boundary layer during accelerated flow events. Geophys. Res. Lett. **17**(11), 1833 (1990a)
- J.T. Gosling, R.M. Skoug, D.J. McComas et al., Magnetic disconnection from the Sun: observations of a reconnection exhaust in the solar wind at the heliospheric current sheet. Geophys. Res. Lett. 32, L05105 (2005). doi:10.1029/2005GL022406
- A. Grocott, S.W.H. Cowley, J.B. Sigwarth, Ionospheric flow during extended intervals of northward but BYdominated IMF. Ann. Geophys. 21, 509 (2003)
- A. Grocott et al., Magnetosonic Mach number dependence of the efficiency of reconnection between planetary and interplanetary magnetic fields. J. Geophys. Res. 114, A07219 (2009). doi:10.1029/2009JA014330
- K.C. Hansen et al., Global MHD simulations of Saturn's magnetosphere at the time of Cassini approach. Geophys. Res. Lett. 32, L20S06 (2005). doi:10.1029/2005GL022835
- M. Harel et al., Quantitative simulations of a magnetospheric substorm, 1. Model logic and overview. J. Geophys. Res. 86, 2217–2241 (1981)
- R.A. Haring et al., Reactive sputtering of simple condensed gases by keV ions. III. Kinetic energy distributions. Nucl. Instrum. Methods B 5, 483 (1984)
- E.G. Harris, On a plasma sheath separating regions of oppositely directed magnetic fields. Nuovo Cimento 23, 115 (1962)
- H. Hasegawa et al., Transport of solar wind into Earth's magnetosphere through rolled-up Kelvin-Helmholtz vortices. Nature 430(7001), 755 (2004). doi:10.1038/nature02799
- H. Hasegawa et al., Kelvin-Helmholtz waves at the Earth's magnetopause: Multiscale development and associated reconnection. J. Geophys. Res. 114, A12207 (2009)

- W.J. Heikkila, R.J. Pellinen, Localized induced electric field within the magnetotail. J. Geophys. Res. 82, 1610 (1977)
- S. Hess, P. Zarka, F. Mottez, Io-Jupiter interaction, millisecond bursts and field aligned potentials. Planet. Space Sci. 55, 89 (2007)
- S. Hess et al., Electric potential jumps in the Io-Jupiter Flux tube. Planet. Space Sci. 57, 23 (2009)
- T.W. Hill, F.C. Michel, Heavy ions from the Galilean satellites and the centrifugal distortion of the Jovian magnetosphere. J. Geophys. Res. 81, 4561 (1976)
- T.W. Hill, A.J. Dessler, F.C. Michel, Configuration of the Jovian magnetosphere. Geophys. Res. Lett. 1 (1974). doi:10.1029/GL001i001p00003
- T.W. Hill et al., Evidence for rotationally-driven plasma transport in Saturn's magnetosphere. Geophys. Res. Lett. 32, L41S10 (2005)
- R.R. Hodges Jr., Monte Carlo simulation of the terrestrial hydrogen exosphere. J. Geophys. Res. 99, 23,229 (1994). doi:10.1029/94JA02183
- W.O. Hofer, Angular, energy, and mass distribution of sputtered particles, in *Sputtering by Particle Bombardment*, ed. by R. Behrisch, K. Wittmaack (Springer, Berlin, 1991), p. 15
- E.W. Hones Jr., The magnetotail: its generation and dissipation, in *Physics of Solar Planetary Environments*, ed. by D.J. Williams (AGU, Washington, 1976), pp. 559–571
- E.W. Hones Jr., Substorm processes in the magnetotail: comments on "On hot tenuous plasma, fireballs, and boundary layers in the Earth's magnetotail" by L.A. Frank et al. J. Geophys. Res. **82**, 5633 (1977)
- T.S. Huang, T.J. Birmingham, The polarization electric field and its effects in an anisotropic, rotating magnetospheric plasma. J. Geophys. Res. 97, 1511 (1992)
- J.D. Huba, Numerical Methods: Ideal and Hall MHD 7, 26 (2005)
- R.L. Huff et al., Mapping of auroral kilometric radiation sources to the Aurora. J. Geophys. Res. 93, 11445 (1988)
- A. Ieda et al., Statistical analysis of the plasmoid evolution with Geotail observations. J. Geophys. Res. 103(A3), 4453 (1998). doi:10.1029/97JA03240
- R. Ilie, R.M. Skoug, H.O. Funsten, M.W. Liemohn, J.J. Bailey, M. Gruntman, The impact of geocoronal density on ring current development. J. Atmos. Sol.-Terr. Phys. 99, 92 (2013). doi:10.1016/j.jastp. 2012.03.010
- S.M. Imber et al., MESSENGER observations of large dayside flux transfer events: do they drive Mercury's substorm cycle? J. Geophys. Res. (2014, submitted)
- F.M. Ipavich et al., Energetic (greater than 100 keV) O⁽⁺⁾ ions in the plasma sheet. Geophys. Res. Lett. 11, 504 (1984)
- C.M. Jackman et al., Interplanetary magnetic field at ~9 AU during the declining phase of the solar cycle and its implications for Saturn's magnetospheric dynamics. J. Geophys. Res. 109, A11203 (2004). doi:10.1029/2004JA010614
- C.M. Jackman et al., Strong rapid dipolarizations in Saturn's magnetotail: In situ evidence of reconnection. Geophys. Res. Lett. 34(11), L11203 (2007)
- C.M. Jackman, R.J. Forsyth, M.K. Dougherty, The overall configuration of the interplanetary magnetic field upstream of Saturn as revealed by Cassini observations. J. Geophys. Res. 113, A08114 (2008). doi:10.1029/2008JA013083
- C.M. Jackman, J.A. Slavin, S.W.H. Cowley, Cassini observations of plasmoid structure and dynamics: Implications for the role of magnetic reconnection in magnetospheric circulation at Saturn. J. Geophys. Res. 116, A10212 (2011). doi:10.1029/2011JA016682
- C.M. Jackman et al., Large-scale structure and dynamics of the magnetotails of Mercury, Earth, Jupiter and Saturn. Space Sci. Rev. 182(1), 85–154. (2014a). doi:10.1007/s11214-014-0060-8
- C.M. Jackman et al., Saturn's dynamic magnetotail: A comprehensive magnetic field and plasma survey of plasmoids and travelling compression regions, and their role in global magnetospheric dynamics. J. Geophys. Res. 119, 5465–5494 (2014b). doi:10.1002/2013JA019388
- X. Jia, M.G. Kivelson, Driving Saturn's magnetospheric periodicities from the atmosphere/ionosphere: Magnetotail response to dual sources. J. Geophys. Res. 117, A11219 (2012). doi:10.1029/2012JA018183
- X.Z. Jia et al., Properties of Ganymede's magnetosphere inferred from improved three-dimensional MHD simulations. J. Geophys. Res. 114, A09209 (2009). doi:10.1029/2009JA014375
- X. Jia et al., Magnetospheric configuration and dynamics of Saturn's magnetosphere: A global MHD simulation. J. Geophys. Res. 117, A05225 (2012b). doi:10.1029/2012JA017575
- X. Jia, M.G. Kivelson, T.I. Gombosi, Driving Saturn's magnetospheric periodicities from the upper atmosphere/ionosphere. J. Geophys. Res. 117, A04215 (2012a). doi:10.1029/2011JA017367
- R.E. Johnson, Energetic Charged-Particle Interactions with Atmospheres and Surfaces, vol. X. Phys. Chem. Space (Springer, Berlin, Heidelberg, New York, 1990), p. 19
- R.E. Johnson, Sputtering and desorption from icy surfaces, in *Solar System Ices*, ed. by B. Schmitt, C. de Bergh (Kluwer Acad., Dordrecht, 1998), p. 303

- R.E. Johnson, Surface chemistry in the Jovian magnetosphere radiation environment, in *Chemical Dynamics in Extreme Environments*, ed. by R. Dessler. Adv. Ser. Phys. Chem., vol. 11 (World Scientific, Singapore, 2001), p. 390, Chap. 8
- R.E. Johnson et al., Charged particle erosion of frozen volatiles in ice grains and comets. Astron. Astrophys. 123, 343 (1983)
- R.E. Johnson, F. Leblanc, B.V. Yakshinskiy, T.E. Madey, Energy distributions for desorption of sodium and potassium from ice: the Na/K ratio at Europa. Icarus 156, 136 (2002)
- R.E. Johnson, M.H. Burger, T.A. Cassidy, F. Leblanc, M. Marconi, W.H. Smyth, Composition and detection of Europa's sputter-induced atmosphere, in *Europa*, ed. by R.T. Pappalardo, W.B. McKinnon, K. Khurana (The University of Arizona Press, Tucson, 2009), p. 507
- M. Jorgensen, M.G. Henderson, E.C. Roelof, G.D. Reeves, H.E. Spence, Charge exchange contribution to the decay of the ring current, measured by energetic neutral atoms (ENAs). J. Geophys. Res. 106, 1931 (2001). doi:10.1029/2000JA000124
- K. Kabin et al., Interaction of Mercury with the solar wind. Icarus 143(2), 397 (2000)
- K. Keika, M. Nose, K. Takahashi, S. Ohtani, P.C. Brandt, D.G. Mitchell, S.P. Christon, R.W. McEntire, Contribution of ion flowout and charge exchange processes to the decay of the storm-time ring current. AGU Fall Meeting Abstracts, p. A565 (2003)
- K. Keika, M. Nose, P.C. Brandt, S. Ohtani, D.G. Mitchell, E.C. Roelof, Contribution of charge exchange loss to the storm time ring current decay: IMAGE/HENA observations. J. Geophys. Res. 111, 11 (2006). doi:10.1029/2006JA011789
- A. Keiling et al., Transient ion beamlet injections into spatially separated PSBL flux tubes observed by Cluster-CIS. Geophys. Res. Lett. 31 (2004)
- K.K. Khurana et al., The origin of Ganymede's polar caps. Icarus 191, 193 (2007)
- A. Kidder, R.M. Winglee, E.M. Harnett, Regulation of the centrifugal interchange cycle in Saturn's inner magnetosphere. J. Geophys. Res. 114, A02205 (2009). doi:10.1029/2008JA013100
- R.M. Killen, W.H. Ip, The surface-bounded atmospheres of mercury and the moon. Rev. Geophys. 37(3), 361 (1999)
- R. Killen et al., Processes that promote and deplete the exosphere of Mercury. Space Sci. Rev. **132**, 433 (2007)
- Y.H. Kim et al., Hydrocarbon ions in the lower ionosphere of Saturn. J. Geophys. Res. 119, 384 (2014)
- N. Kitamura et al., Photoelectron flows in the polar wind during geomagnetically quiet periods. J. Geophys. Res. 117, A07214 (2012). doi:10.1029/2011JA017459
- M.G. Kivelson et al., Intermittent short-duration magnetic field anomalies in the Io torus: Evidence for plasma interchange? Geophys. Res. Lett. 24, 2127 (1997)
- M.G. Kivelson et al., Ganymede's magnetosphere: Magnetometer overview. J. Geophys. Res. 103, 19963 (1998)
- M.G. Kivelson et al., Moon-magnetosphere interaction: a tutorial. Adv. Space Res. 33, 2061 (2004)
- S. Knight, Parallel electric fields. Planet. Space Sci. 21, 741 (1973)
- J.U. Kozyra, M.W. Liemohn, C.R. Clauer, A.J. Ridley, M.F. Thomsen, J.E. Borovsky, J.L. Roeder, V.K. Jordanova, W.D. Gonzalez, Multistep Dst development and ring current composition changes during the 4–6 June 1991 magnetic storm. J. Geophys. Res. 107, 1224 (2002). doi:10.1029/2001JA000023
- N.A. Krall, A.W. Trivelpiece, *Principles of Plasma Physics* (San Francisco Press, Incorporated, San Francisco, 1986)
- S.M. Krimigis et al., Dynamics of Saturn's magnetosphere from MIMI during Cassini's orbital insertion. Science 307, 1270 (2005)
- E.A. Kronberg et al., Comparison of periodic substorms at Jupiter and Earth. J. Geophys. Res. 113, A04212 (2008). doi:10.1029/2007JA012880
- N.A. Krupp et al., Global flows of energetic ions in Jupiter's equatorial plane: First-order approximation. J. Geophys. Res. 106, 26,017 (2001). doi:10.1029/2000JA900138
- N.A. Krupp et al., Dynamics of the Jovian magnetosphere, in *Jupiter: The Planet, Satellites, and Magnetosphere*, ed. by F. Bagenal et al. (Cambridge Univ. Press, Cambridge, 2004), p. 617, Chap. 25
- A. Lagg et al., In situ observations of a neutral gas torus at Europa. Geophys. Res. Lett. **30**, 110000 (2003)
- H.R. Lai et al., Reconnection at the magnetopause of Saturn: Perspective from FTE occurrence and magnetosphere size. J. Geophys. Res. 117, A05222 (2012). doi:10.1029/2011JA017263
- B. Lavraud, E. Larroque, E. Budnik et al., Asymmetry of magnetosheath flows and magnetopause shape during low Alfven Mach number solar wind. J. Geophys. Res. 118, 1089 (2013). doi:10.1002/jgra.50145
- S.A. Ledvina, Y.-J. Ma, E. Kallio, Modeling and simulating flowing plasmas and related phenomena. Space Sci. Rev. 139, 143189 (2008). doi:10.1007/s11214-008-9384-6
- M.W. Liemohn, J.U. Kozyra, Testing the hypothesis that charge exchange can cause a two-phase decay, in *The Inner Magnetosphere: Physics and Modeling*, ed. by T.I. Pulkkinen, N.A. Tsyganenko, R.H.W. Friedel, vol. 155 (American Geophysical Union Geophysical Monograph Series, Washington, 2005), p. 211

A. Lipatov, The Hybrid Multiscale Simulation Technology (Springer, Berlin, 2002)

- P. Louarn et al., Trapped electrons as a free energy source for auroral kilometric radiation. J. Geophys. Res. 95, 5983 (1990)
- J.G. Luhmann et al., A model of the ionosphere of Saturn's rings and its implications. Icarus 181, 465 (2006)
- J.G. Lyon, J.A. Fedder, C.M. Mobarry, The Lyon-Fedder-Mobarry (LFM) global MHD magnetospheric simulation code. J. Atmos. Sol.-Terr. Phys. 66, 1333 (2004). doi:10.1016/j.jastp.2004.03.020
- L.R. Lyons, Generation of large-scale regions of auroral currents, electric potentials, and precipitation by the divergence of the convection electric field. J. Geophys. Res. **85**, 17 (1980)
- R. Lysak, Electrodynamic couplig of the ionosphere and magnetosphere. Space Sci. Rev. 52, 33 (1990)
- T. Majeed, J.C. McConnell, Voyager electron density measurements on Saturn: Analysis with a time dependent ionospheric model. J. Geophys. Res. 101, 7589 (1996). doi:10.1029/96JE00115
- V. Mangano et al., The contribution of impact-generated vapour to the hermean atmosphere. Planet. Space Sci. 55(11), 1541 (2007)
- M.L. Marconi, A kinetic model of Ganymede's atmosphere. Icarus 190, 155 (2007)
- G.T. Marklund, Electric fields and plasma processes in the auroral downward current region, below, within, and above the acceleration region. Space Sci. Rev. **142**, 1 (2009). doi:10.1007/s11214-008-9373-9
- A. Masters et al., Surface waves on Saturn's dawn flank magnetopause driven by the Kelvin-Helmholtz instability. Planet. Space Sci. 57, 1769 (2009). doi:10.1016/j.pss.2009.02.010
- A. Masters et al., Cassini observations of a Kelvin-Helmholtz vortex in Saturn's outer magnetosphere. J. Geophys. Res. 115(A7), A07225 (2010). doi:10.1029/2010JA015351
- A. Masters et al., The importance of plasma b conditions for magnetic reconnection at Saturn's magnetopause. Geophys. Res. Lett. 39, L08103 (2012). doi:10.1029/2012GL051372
- Y. Matsumoto, M. Hoshino, Turbulent mixing and transport of collisionless plasmas across a stratified velocity shear layer. J. Geophys. Res. 111, A05213 (2006). doi:10.1029/2004JA010988
- Y. Matsumoto, K. Seki, Formation of a broad plasma turbulent layer by forward and inverse energy cascades of the Kelvin–Helmholtz instability. J. Geophys. Res. 115, A10231 (2010). doi:10.1029/2009JA014637
- B.H. Mauk, Quantitative modeling of the "convection surge" mechanism of ion acceleration. J. Geophys. Res. 91, 13,423 (1986)
- B.H. Mauk et al., Energetic neutral atoms from a trans-Europa gas torus at Jupiter. Nature 421, 920 (2003)
- H.J. McAndrews et al., Evidence for reconnection at Saturn's magnetopause. J. Geophys. Res. 113, A04210 (2008). doi:10.1029/2007JA012581
- M.B. McElroy, The ionospheres of the major planets. Space Sci. Rev. 14, 460 (1973)
- J. McFadden, C. Carlson, R. Ergun, Microstructure of the auroral acceleration region as observed by FAST. J. Geophys. Res. 104(A7), 14453 (1999). doi:10.1029/1998JA900167
- R.L. McNutt Jr. et al., Departure from rigid co-rotation of plasma in Jupiter's dayside magnetosphere. Nature 280, 803 (1979)
- B. Meinel, Doppler-shifted auroral hydrogen emission. Astrophys. J. 113, 50 (1951). doi:10.1086/145375
- N. Meyer-Vernet, M. Moncuquet, S. Hoang, Temperature inversion in the Io plasma torus. Icarus 116, 202 (1995)
- F.C. Michel, P.A. Sturrock, Centrifugal instability of the Jovian magnetosphere and its interaction with the solar wind. Planet. Space Sci. 22, 1501 (1974)
- D.G. Mitchell et al., Current carriers in the near-earth cross-tail current sheet during substorm growth phase. Geophys. Res. Lett. 17, 583–586 (1990)
- D.G. Mitchell et al., Global imaging of O⁺ from IMAGE HENA. Space Sci. Rev. 109, 63 (2003)
- T.E. Moore, J.L. Horwitz, Stellar ablation of planetary atmospheres. Rev. Geophys. 45, RG3002 (2007). doi:10.1029/2005RG000194
- F. Mottez, V. Génot, Electron acceleration by an Alfvénic pulse propagating in an auroral plasma cavity. J. Geophys. Res. 116, A00K15 (2011)
- F.S. Mozer, A. Hull, Origin and geometry of upward parallel electric fields in the auroral acceleration region. J. Geophys. Res. 106, 5763 (2001)
- F.S. Mozer et al., Observations of paired electrostatic shocks in the polar magnetosphere. Phys. Rev. Lett. **38**, 292 (1977)
- F.S. Mozer et al., Satellite measurements and theories of low altitude auroral particle acceleration. Space Sci. Rev. 27, 155 (1980)
- M. Müller et al., Estimation of the dust flux near Mercury. Planet. Space Sci. 50, 1101 (2002)
- A. Mura et al., The sodium exosphere of Mercury: Comparison between observations during Mercury's transit and model results. Icarus 200, 1 (2009)
- A.F. Nagy, T.E. Cravens, S.G. Smith, H.A. Taylor Jr., H.C. Brinton, Model calculations of the dayside ionosphere of Venus: Ionic composition. J. Geophys. Res. 85, 7795–7801 (1980)
- T.K.M. Nakamura, M. Fujimoto, A. Otto, Structure of an MHD-scale Kelvin–Helmholtz vortex: Twodimensional two-fluid simulations including finite electron inertial effects. J. Geophys. Res. 113, A09204 (2008). doi:10.1029/2007JA012803

- A.I. Neishtadt, On the change in the adiabatic invariant on crossing a separatrix in systems with two degrees of freedom. J. Appl. Math. 51, 586 (1987)
- P.T. Newell et al., Characterizing the state of the magnetosphere: Testing the ion precipitation maxima latitude (b2i) and the ion isotropy boundary. J. Geophys. Res. 103, 4739 (1998)
- J.D. Nichols, S.W.H. Cowley, D.J. McComas, Magnetopause reconnection rate estimates for Jupiter's magnetosphere based on interplanetary measurements at ~5AU. Ann. Geophys. 24, 393 (2006)
- P. Norqvist et al., Ion cyclotron heating in the dayside magnetosphere. J. Geophys. Res. 101, 13,179 (1996)
- T.G. Northrop, The Adiabatic Motion of Charged Particles (Wiley Interscience, New York, 1963)
- T.G. Northrop, J.R. Hill, Stability of negatively charged dust grains in Saturn's ring plane. J. Geophys. Res. 87, 6045 (1983)
- M. Nosé et al., Acceleration of oxygen ions of ionospheric origin in the near-Earth magnetotail during substorms. J. Geophys. Res. 105, 7669 (2000)
- K. Nykyri, A. Otto, Plasma transport at the magnetospheric boundary due to reconnection in Kelvin– Helmholtz vortices. Geophys. Res. Lett. 28(18), 3565 (2001). doi:10.1029/2001GL013239
- K. Nykyri et al., Cluster observations of reconnection due to the Kelvin-Helmholtz instability at the dawnside magnetospheric flank. Ann. Geophys. 24, 2619 (2006)
- Y. Ogawa, K. Seki, M. Hirahara et al., Coordinated EISCAT Svalbard radar and Reimei satellite observations of ion upflows and suprathermal ions. J. Geophys. Res. 113, A05306 (2008). doi:10.1029/ 2007JA012791
- T.R. Ogino, R.J. Walker, M.G. Kivelson, A global magnetohydrodynamic simulation of the Jovian magnetosphere. J. Geophys. Res. 84, 47 (1998)
- M. Øieroset et al., A statistical study of ion beams and conics from the dayside ionosphere during different phases of a substorm. J. Geophys. Res. 104, 6987 (1999)
- Y.M. Ono et al., The role of magnetic field fluctuations in nonadiabatic acceleration of ions during dipolarization. J. Geophys. Res. 114 (2009). doi:10.1029/2008JA013918
- S. Orsini, A. Milillo, Magnetospheric plasma loss processes and energetic neutral atoms. Il Nuovo Cimento 22(5), 633 (1999)
- N. Østgaard, S.B. Mende, H.U. Frey, G.R. Gladstone, H. Lauche, Neutral hydrogen density profiles derived from geocoronal imaging. J. Geophys. Res. 108, 1300 (2003). doi:10.1029/2002JA009749
- L. Pallier, R. Prangé, Detection of the southern counterpart of the north FUV polar cusp. Shared properties. Geophys. Res. Lett. 31, L06701 (2004)
- C. Paranicas et al., The ion environment near Europa and its role in surface energetics. Geophys. Res. Lett. 29(5), 1074 (2002). doi:10.1029/2001GL014127
- E.N. Parker, Dynamics of the interplanetary magnetic field. Astrophys. J. 128, 664 (1958)
- G.S. Paschmann, Recent in-situ observations of magnetic reconnection in near-Earth space. Geophys. Res. Lett. 35, L19109 (2008). doi:10.1029/2008GL035297
- G. Paschmann, S. Haaland, R. Treumann (eds.), Auroral Plasma Physics (Kluwer Academic, Dordrecht, 2003)
- G. Paschmann, M. Øieroset, T. Phan, In-situ observations of reconnection in space. Space Sci. Rev. 178, 385–417 (2013). doi:10.1007/s11214-012-9957-2
- R.J. Pellinen, W.J. Heikkila, Energization of charged particles to high energies by an induced substorm electric field within the magnetotail. J. Geophys. Res. 83, 1544 (1978)
- P. Perreault, S.I. Akasofu, Study of geomagnetic storms. Geophys. J. R. Astron. Soc. 54(3), 547 (1978)
- C. Peymirat, D. Fontaine, Numerical simulation of the magnetospheric convection including the effect of field-aligned currents and electron precipitation. J. Geophys. Res. 99, 11155 (1994)
- T.D. Phan, M. Oieroset, M. Fujimoto, Reconnection at the dayside low-latitude magnetopause and its nonrole in low-latitude boundary layer formation during northward interplanetary magnetic field. Geophys. Res. Lett. 32, L17101 (2005). doi:10.1029/2005GL023355
- T.D. Phan, G. Paschmann, J.T. Gosling, M. Oieroset, M. Fujimoto, J.F. Drake, V. Angelopoulos, The dependence of magnetic reconnection on plasma b and magnetic shear: evidence from magnetopause observations. Geophys. Res. Lett. 40, 11 (2013). doi:10.1029/2012GL054528
- W.G. Pilipp, G. Morfill, The formation of the plasma sheet resulting from plasma mantle dynamics. J. Geophys. Res. 83, 5670 (1978)
- C. Plainaki et al., Neutral particle release from Europa's surface. Icarus 210, 385 (2010)
- C. Plainaki et al., The role of sputtering and radiolysis in the generation of Europa exosphere. Icarus 218(2), 956 (2012). doi:10.1016/j.icarus.2012.01.023
- C. Plainaki et al., Exospheric O₂ densities at Europa during different orbital phases. Planet. Space Sci. 88, 42 (2013)
- C. Plainaki et al., The H₂O and O₂ exospheres of Ganymede: the result of a complex interaction between the jovian magnetospheric ions and the icy moon. Icarus **245**, 306 (2015)
- D.H. Pontius, Coriolis influences on the interchange instability. Geophys. Res. Lett. 24, 2961 (1997)

- A.E. Potter, Chemical sputtering could produce sodium vapour and ice on Mercury. Geophys. Res. Lett. 22(23), 3289 (1995)
- K.G. Powell et al., A solution-adaptive upwind scheme for ideal magnetohydrodynamics. J. Comput. Phys. 154(2), 284 (1999). doi:10.1006/jcph.1999.6299
- R. Prangé, The UV and IR Jovian aurorae. Adv. Space Res. 12(8), 379 (1992)
- R. Prangé et al., Rapid energy dissipation and variability of the Io-Jupiter electrodynamic circuit. Nature 379, 323 (1996)
- R. Prangé et al., Detailed study of FUV Jovian auroral features with the post COSTAR Hubble Faint Object Camera. J. Geophys. Res. 103, 20195 (1998)
- P.L. Pritchett, Electron-cyclotron maser instability in relativistic plasmas. Phys. Fluids 29, 2919 (1986)
- W.R. Pryor et al., The auroral footprint of Enceladus on Saturn. Nature 472, 331 (2011)
- J.D. Raeder et al., Open GGCM simulations for the THEMIS mission. Space Sci. Rev. 141, 535 (2008). doi:10.1007/s11214-008-9421-5
- J. Raines et al., Space Sci. Rev. (2015, this issue)
- R.L. Rairden, L.A. Frank, J.D. Craven, Geocoronal imaging with Dynamics Explorer. J. Geophys. Res. 91, 13,613 (1986). doi:10.1029/JA091iA12p13613
- L.C. Ray et al., Current-voltage relation of a centrifugally confined plasma. J. Geophys. Res. **114**, A04214 (2009). doi:10.1029/2008JA013969
- C.T. Reimann et al., Ion-induced molecular ejection from D₂O ice. Surf. Sci. 147, 227 (1984)
- H.E. Roosendaal, R.A. Hating, J.B. Sanders, Surface disruption as an observable factor in the energy distribution of sputtered particles. Nucl. Instrum. Methods 194, 579 (1982)
- H. Rosenbauer, H. Grunwaldt, M.D. Montgomery, G. Paschmann, N. Sckopke, Heos 2 plasma observations in the distant polar magnetosphere: the plasma mantle. J. Geophys. Res. 80, 2723 (1975)
- A.A. Roux et al., Auroral kilometric radiation sources: in situ and remote sensing observations from Viking. J. Geophys. Res. 98, 11657 (1993)
- C.T. Russell et al., Localized reconnection in the near Jovian magnetotail. Science 280, 1061 (1998). doi: 10.1126/science.280.5366.1061
- P.E. Sandholt et al., Dayside auroral configurations: Responses to southward and northward rotations of the interplanetary magnetic field. J. Geophys. Res. **103**(20), 279 (1998)
- R.W. Schunk, A.F. Nagy, Ionospheres of the terrestrial planets. Rev. Geophys. 18, 813-852 (1980)
- R.W. Schunk, A.F. Nagy, *Ionospheres*, 2nd edn. (Cambridge University Press, Cambridge, 2009)
- R.W. Schunk, J.J. Sojka, Global ionosphere–polar wind system during changing magnetic activity. J. Geophys. Res. 102, 11625 (1997)
- L. Scurry, C.T. Russell, Proxy studies of energy transfer to the magnetosphere. J. Geophys. Res. 96, 9541 (1991)
- V.A. Sergeev et al., Pitch-angle scattering of energetic protons in the magnetotail current sheet as the dominant source of their isotropic precipitation into the nightside ionosphere. Planet. Space Sci. 31, 1147 (1983)
- V.A. Sergeev, M. Malkov, K. Mursula, Testing the isotropic boundary algorithm to evaluate the magnetic field configuration of the tail. J. Geophys. Res. 98, 7609 (1993)
- V.P. Shabansky, Some processes in the magnetosphere. Space Sci. Rev. 12, 299 (1971)
- V.I. Shematovich et al., Surface-bounded atmosphere of Europa. Icarus 173, 480 (2005)
- M. Shi, R.A. Baragiola, D.E. Grosjean, R.E. Johnson, S. Jurac, J. Schou, Sputtering of water ice surfaces and the production of extended neutral atmospheres. J. Geophys. Res. 100, 26387 (1995)
- P. Sigmund, Theory of Sputtering. I. Sputtering Yield of Amorphous and Polycrystalline (1969)
- G.L. Siscoe, On the equatorial confinement and velocity space distribution of satellite ions in Jupiter's magnetosphere. J. Geophys. Res. 82, 1641 (1977)
- G. Siscoe, L. Christopher, Variations in the solar wind stand-off distance at Mercury. Geophys. Res. Lett. 2, 158 (1975)
- J.A. Slavin, Traveling compression regions, in *New Perspectives in Magnetotail Physics*, ed. by A. Nishida, S.W.H. Cowley, D.N. Baker. AGU Monograph, vol. 105 (AGU, Washington, 1998), pp. 225–240
- J.A. Slavin, R.E. Holzer, J.R. Spreiter, S.S. Stahara, Planetary mach cones: theory and observation. J. Geophys. Res. 89, 2708 (1984)
- J.A. Slavin et al., Geotail observations of magnetic flux ropes in the plasma sheet. J. Geophys. Res. 108(A1), 1015 (2003). doi:10.1029/2002JA009557
- J.A. Slavin, M.H. Acuna, B.J. Anderson et al., MESSENGER observations of magnetic reconnection in Mercury's magnetosphere. Science 324(5927), 606 (2009). doi:10.1126/science.1172011
- J.A. Slavin et al., MESSENGER and Mariner 10 flyby observations of magnetotail structure and dynamics at Mercury. J. Geophys. Res. 117, A01215 (2012a). doi:10.1029/2011JA016900
- J.A. Slavin et al., MESSENGER observations of flux transfer events at Mercury. J. Geophys. Res. 117, A00M06 (2012b). doi:10.1029/2012JA017926

- J.A. Slavin et al., MESSENGER observations of Mercury's dayside magnetosphere under extreme solar wind conditions. J. Geophys. Res. 119, 8087–8116 (2014)
- P.H. Smith, N.K. Bewtra, Charge exchange lifetimes for ring current ions. Space Sci. Rev. 22, 301 (1978). doi:10.1007/BF00239804
- W.H. Smyth, M.L. Marconi, Europa's atmosphere, gas tori, and magnetospheric implications. Icarus 181, 510 (2006)
- S.C. Solomon, Auroral electron transport using the Monte Carlo method. Geophys. Res. Lett. 20, 185 (1993)
- B.U.O. Sonnerup, Adiabatic particle orbits in a magnetic null sheet. J. Geophys. Res. **76**, 8211 (1971)
- B.U.O. Sonnerup, The magnetopause reconnection rate. J. Geophys. Res. 79, 1546 (1974). doi:10.1029/ JA079i010p01546
- T.W. Speiser, Particle trajectory in model current sheets, 1, Analytical solutions. J. Geophys. Res. **70**, 4219 (1965)
- W.N. Spjeldvik, Equilibrium structure of equatorially mirroring radiation belt protons. J. Geophys. Res. 82, 2801 (1977). doi:10.1029/JA082i019p02801
- W.N. Spjeldvik, T.A. Fritz, Theory for charge states of energetic oxygen ions in the earth's radiation belts. J. Geophys. Res. 83, 1583 (1978). doi:10.1029/JA083iA04p01583
- R.J. Strangeway et al., Factors controlling ionospheric outflows as observed at intermediate altitudes. J. Geophys. Res. 110, A03221 (2005). doi:10.1029/2004JA010829
- G.W. Stuart, Satellite-measured radiation. Phys. Rev. Lett. 2, 417 (1959)
- M. Sugiura, Hourly Values of Equatorial Dst for the IGY. Ann. Int. Geophys. Year, vol. 35 (Pergamon Press, Oxford, 1964), p. 9
- T.S. Sundberg et al., MESSENGER orbital observations of large-amplitude Kelvin-Helmholtz waves at Mercury's magnetopause. J. Geophys. Res. **117**, A04216 (2012)
- K. Szegö et al., Physics of mass loaded plasmas. Space Sci. Rev. 94, 429 (2000)
- S.W.Y. Tam, T. Chang, V. Pierrard, Kinetic modeling of the polar wind. J. Atmos. Sol.-Terr. Phys. 69, 1984 (2007)
- B.D. Teolis, R.A. Vidal, J. Shi, R.A. Baragiola, Mechanisms of O₂ sputtering from water ice by keV ions. Phys. Rev. B 72, 245422 (2005). 2005. doi:10.1103/PhysRevB.72.245422
- N. Terada, H. Shinagawa, T. Tanaka, K. Murawski, K. Terada, A three-dimensional, multispecies, comprehensive MHD model of the solar wind interaction with the planet Venus. J. Geophys. Res. 114, A09208 (2009). doi:10.1029/2008JA013937
- T. Terasawa et al., Solar wind control of density and temperature in the near-Earth plasma sheet: WIND/GEOTAIL collaboration. Geophys. Res. Lett. **24**(8), 935 (1997). doi:10.1029/96GL04018
- B.T. Thomas, E.J. Smith, The Parker spiral configuration of the interplanetary magnetic field between 1 and 8.5 AU. J. Geophys. Res. 85, 6861 (1980)
- N. Thomas, F. Bagenal, T.W. Hill, J.K. Wilson, The Io neutral cloud and plasma torus, in *Jupiter: The Planet, Satellites, and Magnetosphere*, ed. by F. Bagenal et al. (Cambridge Univ. Press, Cambridge, 2004), p. 560, Chap. 23
- M.F. Thomsen et al., Survey of ion plasma parameters in Saturn's magnetosphere. J. Geophys. Res. 115, A10220 (2010). doi:10.1029/2010JA015267
- R.M. Thorne et al., Galileo evidence for rapid interchange transport in the Io torus. Geophys. Res. Lett. 24, 2131 (1997)
- G. Toth et al., Space weather modeling framework: a new tool for the space science community. J. Geophys. Res. 110, 12,226 (2005). doi:10.1029/2005JA011126
- G. Toth et al., Adaptive numerical algorithms in space weather modeling. J. Comput. Phys. 231(3), 870–903 (2012). doi:10.1016/j.jcp.2011.02.006
- R.A. Treumann, The electron–cyclotron maser for astrophysical application. Astron. Astrophys. Rev. 13, 229 (2006)
- V. Vasyliunas, Mathematical models of magnetospheric convection and its coupling to the ionosphere, in Particles and Fields in the Magnetosphere, ed. by B. McCormac (Reidel, Hingham, 1970)
- V.M. Vasyliunas, Plasma distribution and flow, in *Physics of the Jovian Magnetosphere*, ed. by A.J. Dessler (Cambridge Univ. Press, Cambridge, 1983), p. 395
- M.F. Vogt et al., Reconnection and flows in the Jovian magnetotail as inferred from magnetometer observations. J. Geophys. Res. 115, A06219 (2010). doi:10.1029/2009JA015098
- M.F. Vogt et al., Structure and statistical properties of plasmoids in Jupiter's magnetotail. J. Geophys. Res. 119, 821 (2014). doi:10.1002/2013JA019393
- J.H. Waite et al., An auroral flare at Jupiter. Nature 410, 787 (2001)
- J.A. Wanliss, K.M. Showalter, High-resolution global storm index: Dst versus SYM-H. J. Geophys. Res. 111(A2), A02202 (2006). doi:10.1029/2005JA011034
- D. Welling et al., Space Sci. Rev. (2015, this issue)

- M.S. Westley et al., Photodesorption from low-temperature water ice in interstellar and circumstellar grains. Nature 373, 405 (1995)
- D.J. Williams, Ring current composition and sources: An update. Planet. Space Sci. 29, 1195 (1981)
- R.J. Wilson, R.L. Tokar, M.G. Henderson, Thermal ion flow in Saturn's inner magnetosphere measured by the Cassini plasma spectrometer: A signature of the Enceladus torus? Geophys. Res. Lett. 36, L23104 (2009). doi:10.1029/2009GRL040225
- M. Wiltberger et al., Influence of cusp O⁺ outflow on magnetotail dynamics in a multifluid MHD model of the magnetosphere. J. Geophys. Res. 115, A00J05 (2010). doi:10.1029/2010JA015579
- S. Wing, P.T. Newell, 2D plasma sheet ion density and temperature profiles for northward and southward IMF. Geophys. Res. Lett. 29(9), 1307 (2002). doi:10.1029/2001GL013950
- R.M. Winglee et al., Global impact of ionospheric outflows on the dynamics of the magnetosphere and crosspolar cap potential. J. Geophys. Res. 107, 1237 (2002). doi:10.1029/2001JA000214
- D. Winske, L. Yin, N. Omidi, H. Karimabadi, K. Quest, Hybrid simulation codes: Past, present and future— A tutorial, in *Space Plasma Simulation*, ed. by J. Büchner, C. Dum, M. Scholer. Lect. Notes Phys., vol. 615 (2003), pp. 136–165
- R.A. Wolf, Ionosphere-magnetosphere coupling. Space Sci. Rev. 17, 535 (1979)
- C.S. Wu, L.C. Lee, A theory of the terrestrial kilometric radiation. Astrophys. J. 230, 621 (1979)
- X.-Y. Wu, J.L. Horwitz, J.-N. Tu, Dynamic fluid kinetic (DyFK) simulation of auroral ion transport: Synergistic effects of parallel potentials, transverse ion heating, and soft electron precipitation. J. Geophys. Res. 107(A10), 1283 (2002). doi:10.1029/2000JA000190
- P. Wurz, H. Lammer, Monte-Carlo simulation of Mercury's exosphere. Icarus 164, 1 (2003)
- P. Wurz et al., The lunar exosphere: the sputtering contribution. Icarus 191, 486–496 (2007). doi:10.1016/ j.icarus.2007.04.034
- P. Wurz et al., Self-consistent modelling of Mercury's exosphere by sputtering, micro-meteorite impact and photon-stimulated desorption. Planet. Space Sci. 58, 1599 (2010)
- B.V. Yakshinskiy, T.E. Madey, Photon-stimulated desorption as a substantial source of sodium in the lunar atmosphere. Nature **400**, 642 (1999)
- B.V. Yakshinskiy, T.E. Madey, Electron- and photon-stimulated desorption of K from ice surfaces. J. Geophys. Res. 106, 33303 (2001)
- B.V. Yakshinskiy, T.E. Madey, Temperature-dependent DIET of alkalis from SiO₂ films: Comparison with a lunar sample. Surf. Sci. 593, 202 (2005)
- A.W. Yau, M. Andre, Sources of ion outflow in the high latitude ionosphere. Space Sci. Rev. 80, 1 (1997). doi:10.1023/A:1004947203046
- A.W. Yau, T. Abe, W.K. Peterson, The polar wind: Recent observations. J. Atmos. Sol.-Terr. Phys. 69, 1936 (2007)
- K.S. Yee, Numerical solution of initial boundary value problems involving Maxwell's equations in isotropic media. IEEE Trans. Antennas Propag. 14, 302 (1966)
- Y. Yu, A.J. Ridley, Exploring the influence of ionospheric O⁺ outflow on magnetospheric dynamics: dependence on the source location. J. Geophys. Res. 118, 1711 (2013). doi:10.1029/2012JA018411
- P. Zarka, Auroral radio emissions at the outer planets: observations and theories. J. Geophys. Res. 103, 20159 (1998)
- J.F. Ziegler, SRIM-2003. Nucl. Instrum. Methods B 219, 1027 (2004)
- J.F. Ziegler, J.P. Biersack, U. Littmark, *The Stopping and Range of Ions in Solids*. Stopping and Ranges of Ions in Matter, vol. 1 (Pergamon Press, New York, 1984)
- J.H. Zoennchen, U. Nass, G. Lay, H.J. Fahr, 3-D-geocoronal hydrogen density derived from TWINS Lyalpha-data. Ann. Geophys. 28, 1221 (2010). doi:10.5194/angeo-28-1221-2010
- J.H. Zoennchen, J.J. Bailey, U. Nass, M. Gruntman, H.J. Fahr, J. Goldstein, The TWINS exospheric neutral H-density distribution under solar minimum conditions. Ann. Geophys. 29, 2211 (2011)
- B.J. Zwan, R.A. Wolf, Depletion of solar-wind plasma near a planetary boundary. J. Geophys. Res. 81, 1636 (1976)